

第十二届新物理研讨会 青岛 2023.07



LLP scalar searches at FASER

Wei Su

2212.06186 (F. Kling, S. Li, S. Su, H.Song, WS)


outline

- Motivation: Brief introduction to LLP
- Method: Brief introduction to FASER
- General study
 - Production
 - Decay
 - Constraints
- Case study: 2HDM results

Motivation:LLP

mass →	≈2.3 MeV/c ²	≈1.275 GeV/c ²	≈173.07 GeV/c ²	0	≈126 GeV/c ²
charge →	2/3	2/3	2/3	0	0
spin →	1/2	1/2	1/2	1	0
	u up	c charm	t top	g gluon	H Higgs boson
QUARKS					
	≈4.8 MeV/c ²	≈95 MeV/c ²	≈4.18 GeV/c ²	0	
	-1/3	-1/3	-1/3	0	
	1/2	1/2	1/2	1	
	d down	s strange	b bottom	γ photon	
	0.511 MeV/c ²	105.7 MeV/c ²	1.777 GeV/c ²	91.2 GeV/c ²	
	-1	-1	-1	0	
	1/2	1/2	1/2	1	
	e electron	μ muon	τ tau	Z Z boson	
LEPTONS					
	<2.2 eV/c ²	<0.17 MeV/c ²	<15.5 MeV/c ²	80.4 GeV/c ²	
	0	0	0	±1	
	1/2	1/2	1/2	1	
	ν_e electron neutrino	ν_μ muon neutrino	ν_τ tau neutrino	W W boson	
					GAUGE BOSONS


LLP



Weakly interacting
Light Particle




Strongly interacting
Heavy Particle



Heavy

FASER: ForwArd Search ExpeRiment

$pp \rightarrow \text{LLP} + X$, LLP travels ~ 480 m, $\text{LLP} \rightarrow \text{charged tracks} + X$



many hadrons: 10^{17} π , 10^{16} K , 10^{15} D , 10^{14} B with $E \sim \text{TeV}$

Production

PP \rightarrow mesons \rightarrow LLP + X, LLP \rightarrow charged tracks + X

Data base: FORESEE : [2105.07077](https://arxiv.org/abs/2105.07077) F. Kling and S. Trojanowski,

Particle category	Particles	Generators			
		EPOS-LHC	QGSJET II-04	SIBYLL 2.3c	Pythia 8.2
Photons	γ	✓	✓	✓	
Light hadrons	$\pi^0, \pi^+, \eta, \eta', \omega, \rho, \phi, n, p$ $K^+, K_L, K_S, K_0^*, K^{*+}, \Lambda$	✓	✓	✓	
Charm hadrons	$D^+, D^0, D_s^+, \Lambda_c$			✓	✓
Beauty hadrons	$B^0, B^+, B_s, B_c^+, \Lambda_b$				✓
Heavy quarks	c, b				✓
Quarkonia	$J/\Psi, \psi(2S), \Upsilon(nS)$				✓
Weak bosons	W^+, Z, h				✓

Production

Data base: FORESEE

- the `distance` from the IP in meter (default: 480)
- the `length` in meter (default: 5)
- the `luminosity` in units of fb⁻¹ (default: 3000)
- the `selection` : (default: `np.sqrt(x.x**2 + x.y**2) < 1`)
- the decay `channels` which the detector can see

Particle category	Particles	Generators			
		EPOS-LHC	QGSJET II-04	SIBYLL 2.3c	Pythia 8.2
Photons	γ	✓	✓	✓	
Light hadrons	$\pi^0, \pi^+, \eta, \eta', \omega, \rho, \phi, n, p$ $K^+, K_L, K_S, K_0^*, K^{*+}, \Lambda$	✓	✓	✓	
Charm hadrons	$D^+, D^0, D_s^+, \Lambda_c$			✓	✓
Beauty hadrons	$B^0, B^+, B_s, B_c^+, \Lambda_b$				✓
Heavy quarks	c, b				✓
Quarkonia	$J/\Psi, \psi(2S), \Upsilon(nS)$				✓
Weak bosons	W^+, Z, h				✓

Production

Pion


boosted mesons highly collimated

$$p \cdot \theta = p_T \sim \Lambda_{QCD}$$

$$N_\pi = 10^{18} \text{ at } 3000 \text{ fb}^{-1}$$

FASER: radius $R = 10$ cm,
length $D = 1.5$ m,
FASER 2: radius $R = 1$ m,
length $D = 5$ m.

FORESEE



Production: CP even scalar




$$\mathcal{L} = -\frac{1}{2}m_\phi^2\phi^2 - \sum_f \xi_\phi^f \frac{m_f}{v} \phi \bar{f} f + \xi_\phi^W \frac{2m_W^2}{v} \phi W^{\mu+} W_\mu^- + \xi_\phi^Z \frac{m_Z^2}{v} \phi Z^\mu Z_\mu$$

$$+ \xi_{\phi\phi}^W \frac{g^2}{4} \phi\phi W^{\mu+} W_\mu^- + \xi_{\phi\phi}^Z \frac{g^2}{8 \cos^2 \theta_W} \phi\phi Z^\mu Z_\mu + \xi_\phi^g \frac{\alpha_s}{12\pi v} \phi G_{\mu\nu}^a G^{a\mu\nu} + \xi_\phi^\gamma \frac{\alpha_{ew}}{4\pi v} \phi F_{\mu\nu} F^{\mu\nu}$$

$$K \rightarrow \pi\phi, \eta^{(\prime)} \rightarrow \pi\phi, D \rightarrow X_u\phi, B \rightarrow X_s\phi$$

$$\pi^+ \rightarrow \ell\nu\phi \quad K^+ \rightarrow \ell\nu\phi \quad \Upsilon \rightarrow \phi\gamma$$

meson	quark content	mass (MeV)
π^\pm	$u\bar{d}$	139.57018 ± 0.00035
π^0	$\frac{u\bar{u}-d\bar{d}}{\sqrt{2}}$ [a]	134.9766 ± 0.0006
η	$\frac{u\bar{u}+d\bar{d}-2s\bar{s}}{\sqrt{6}}$ [a]	547.853 ± 0.024
η'	$\frac{u\bar{u}+d\bar{d}+s\bar{s}}{\sqrt{3}}$ [a]	957.66 ± 0.24




Production: CP even scalar

Main contribution

$$K^\pm: 493.677 \pm 0.016 \text{ MeV/}$$


$$K^0: 497.611 \pm 0.013 \text{ MeV/}$$



$$\text{Br}(K^\pm \rightarrow \pi^\pm \phi) = \frac{1}{\Gamma_{K^\pm}} \frac{2p_\phi^0}{m_{K^\pm}} \frac{|\mathcal{M}|^2}{16\pi m_{K^\pm}},$$

$$\mathcal{M}(K^\pm \rightarrow \pi^\pm \phi) = G_F^{1/2} 2^{1/4} \xi_\phi^W \left[\frac{7\lambda(m_{K^\pm}^2 + m_{\pi^\pm}^2 - m_\phi^2)}{18} - \frac{7Am_{K^\pm}^2}{9} \right] + \frac{\xi_\phi^{ds}}{2v} m_s \frac{m_{K^\pm}^2 - m_{\pi^\pm}^2}{m_s - m_d} f_0^{K^\pm \pi^\pm}(q^2)$$

$$\frac{\text{Br}(B \rightarrow X_s \phi)}{\text{Br}(B \rightarrow X_c e \nu)} = \frac{\Gamma(b \rightarrow s \phi)}{\Gamma(b \rightarrow c e \nu)} = \frac{12\pi^2 v^2}{m_b^2} \left(1 - \frac{m_\phi^2}{m_b^2}\right)^2 \frac{1}{f(m_c^2/m_b^2)} \left| \frac{\xi_\phi^{bs}}{V_{cb}} \right|^2$$



b: 4.18 GeV, B: around 5.3 GeV,

Production: CP odd scalar

$$\begin{aligned}
 \mathcal{L}_A = & -\frac{1}{2}m_A^2 A^2 + \sum_{f=u,d,e} \xi_A^f \frac{im_f}{v} \bar{f} \gamma_5 f A + \xi_{AA}^W \frac{g^2}{4} A A W^{\mu+} W_{\mu}^- + \xi_{AA}^Z \frac{g^2}{8 \cos^2 \theta_W} A A Z^{\mu} Z_{\mu} \\
 & + \xi_A^g \frac{\alpha_s}{4\pi v} A G_{\mu\nu}^a \tilde{G}^{a\mu\nu} + \xi_A^{\gamma} \frac{\alpha_{ew}}{4\pi v} A F_{\mu\nu} \tilde{F}^{\mu\nu}, \tag{3.1}
 \end{aligned}$$


CP-odd particle mixing production: contribute mainly at **mass peak**

$$\begin{aligned}
 A &= O_{A3} \pi_3 + O_{A8} \pi_8 + O_{A9} \pi_9 + O_{AA} A \\
 &\approx O_{A3} \pi_3 + O_{A\eta} \eta + O_{A\eta'} \eta' + O_{AA} A
 \end{aligned}$$

$$\mathcal{L}_\chi \ni -\frac{1}{2} (\pi_3 \ \pi_8 \ \pi_9 \ A) \begin{pmatrix} m_\pi^2 & 0 & 0 & \delta m_3^2 \\ 0 & m_{\pi_8}^2 & \Delta & \delta m_8^2 \\ 0 & \Delta & m_{\pi_9}^2 & \delta m_9^2 \\ \delta m_3^2 & \delta m_8^2 & \delta m_9^2 & \bar{m}_A^2 \end{pmatrix} \begin{pmatrix} \pi_3 \\ \pi_8 \\ \pi_9 \\ A \end{pmatrix}$$

$$\sigma_A \approx O_{A\pi^0}^2 \sigma_{\pi^0} + O_{A\eta}^2 \sigma_\eta + O_{A\eta'}^2 \sigma_{\eta'},$$

Decay : CP even scalar



Well studied

$A \rightarrow \gamma\gamma$	$H \rightarrow \gamma\gamma$
$A \rightarrow e^+e^-$	$H \rightarrow e^+e^-$
$A \rightarrow \mu^+\mu^-$	$H \rightarrow \mu^+\mu^-$
$A \rightarrow \tau^+\tau^-$	$H \rightarrow \tau^+\tau^-$
$A \rightarrow q\bar{q}$	$H \rightarrow c\bar{c}$
$A \rightarrow gg$	$H \rightarrow s\bar{s}$
	$H \rightarrow gg$

Scale-ind

Scale > 2/3 GeV

?

$H \rightarrow \pi\pi$

$H \rightarrow KK$

$H \rightarrow \pi\pi\pi\pi$

Chiral Perturbativity...

arXiv:1809.01876

arXiv:1612.06538

- $A \rightarrow \pi\pi\pi$
- $A \rightarrow \eta\pi\pi$
- $A \rightarrow \eta'\pi\pi$
- $A \rightarrow \eta\eta\pi$
- $A \rightarrow KK\pi$
- $A \rightarrow \gamma\pi\pi$
- $A \rightarrow \eta\eta'\pi$
- $A \rightarrow \eta'\eta'\pi$
- $A \rightarrow \eta\eta\eta$
- $A \rightarrow \eta\eta\eta'$
- $A \rightarrow \eta\eta'\eta'$
- $A \rightarrow \eta'\eta'\eta'$
- $A \rightarrow \eta KK$
- $A \rightarrow \eta' KK$

Decay: CP even scalar

$$m_\phi < 2 \text{ GeV}$$

$$H \rightarrow \pi\pi$$

$$H \rightarrow KK$$

$$H \rightarrow \pi\pi\pi\pi$$

$$\Gamma_{\pi\pi} = \frac{3G_F}{16\sqrt{2}\pi m_\Phi} \beta_\pi \left| \xi_\Phi^{gg} \frac{2}{27} (\Theta_\pi - \Gamma_\pi - \Delta_\pi) + \frac{m_u \xi_\Phi^u + m_d \xi_\Phi^d}{m_u + m_d} \Gamma_\pi + (\xi_\Phi^s) \Delta_\pi \right|^2$$

$$\Gamma_{KK} = \frac{G_F}{4\sqrt{2}\pi m_\Phi} \beta_K \left| \xi_\Phi^{gg} \frac{2}{27} (\Theta_K - \Gamma_K - \Delta_K) + \frac{m_u \xi_\Phi^u + m_d \xi_\Phi^d}{m_u + m_d} \Gamma_K + (\xi_\Phi^s) \Delta_K \right|^2$$

$$\Gamma_\pi = \langle \pi\pi | m_u \bar{u}u + m_d \bar{d}d | 0 \rangle, \quad \Delta_\pi = \langle \pi\pi | m_s \bar{s}s | 0 \rangle, \quad \Theta_\pi = \langle \pi\pi | \Theta_\mu^\mu | 0 \rangle$$

Leading order chiral perturbation theory

$$\Gamma_\pi^0 = m_\pi^2, \quad \Delta_\pi^0 = 0, \quad \Theta_\pi^0 = s + 2m_\pi^2$$

$$\Gamma_K^0 = \frac{1}{2} m_\pi^2, \quad \Delta_K^0 = m_K^2 - \frac{1}{2} m_\pi^2, \quad \Theta_K^0 = s + 2m_K^2$$



$$m_\phi < 0.5 \text{ GeV}$$

Decay: CP even scalar

$m_\phi < 2 \text{ GeV}$

$H \rightarrow \pi\pi$
 $H \rightarrow KK$
 $H \rightarrow \pi\pi\pi\pi$

Dispersive Analysis



$m_\phi > 0.5 \text{ GeV}$

$$\Gamma_\pi^0 = m_\pi^2,$$

$$\Delta_\pi^0 = 0,$$

$$\Theta_\pi^0 = s + 2m_\pi^2$$

$$\Gamma_K^0 = \frac{1}{2}m_\pi^2,$$

$$\Delta_K^0 = m_K^2 - \frac{1}{2}m_\pi^2,$$

$$\Theta_K^0 = s + 2m_K^2$$

Decay: CP odd scalar

Follow arXiv:1612.06538

Hadronic decays into Tri-meson for $m_A \lesssim 1.3$ GeV

Radiative Hadronic Decays for $m_A \lesssim 1.3$ GeV

hadronic decays for 1.3 GeV $\lesssim m_A \lesssim 3$ GeV (Spectator Model)

$$A \rightarrow \pi\pi\pi$$

$$A \rightarrow \eta\pi\pi$$

$$A \rightarrow \eta'\pi\pi$$

$$A \rightarrow \eta\eta\pi$$

$$A \rightarrow KK\pi$$

$$A \rightarrow \gamma\pi\pi$$

$$A \rightarrow \eta\eta'\pi$$

$$A \rightarrow \eta'\eta'\pi$$

$$A \rightarrow \eta\eta\eta$$

$$A \rightarrow \eta\eta\eta'$$

$$A \rightarrow \eta\eta'\eta'$$

$$A \rightarrow \eta'\eta'\eta'$$

$$A \rightarrow \eta KK$$

$$A \rightarrow \eta' KK$$

Constraints

- current experiments (typically considered)

SN1987a

LEP $\bar{e}e \rightarrow Z^* \phi$

CHARM *dipoton*

NA62 $K \rightarrow \pi \Phi$

E949 $K^+ \rightarrow \pi^+ + \bar{\nu} \nu$

KTeV $K_L \rightarrow \pi + \bar{\mu} \mu$

LHCb $B \rightarrow K^{(*)} + \bar{\mu} \mu$

MicroBooNE $K^+ \rightarrow \pi^+ \chi (e^+ e^-)$

Case study: 2HDM

- Two Higgs Doublet Model

$$\Phi_i = \begin{pmatrix} \phi_i^+ \\ (v_i + \phi_i^0 + iG_i)/\sqrt{2} \end{pmatrix}$$

$$v_u^2 + v_d^2 = v^2 = (246\text{GeV})^2$$

$$\tan \beta = v_u/v_d$$

	ϕ_1	ϕ_2
Type I	u,d,l	
Type II	u	d,l
lepton-specific	u,d	l
flipped	u,l	d

$$\begin{pmatrix} H^0 \\ h^0 \end{pmatrix} = \begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} \phi_1^0 \\ \phi_2^0 \end{pmatrix},$$

$$A = -G_1 \sin \beta + G_2 \cos \beta$$

$$H^\pm = -\phi_1^\pm \sin \beta + \phi_2^\pm \cos \beta$$

- Parameters (CP-conserving, Flavor Limit, Z_2 Symmetry)

$$m_{11}^2, m_{22}^2, \lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5$$



$$v, \tan \beta, \alpha, m_h, m_H, m_A, m_{H^\pm}$$

Soft Z_2 symmetry breaking: m_{12}^2

246 GeV

125. GeV

Case study: 2HDM

$$\text{Generally: } \cos(\beta - \alpha) = 0$$

	ξ_H^u	ξ_H^d	ξ_H^ℓ	ξ_A^u	ξ_A^d	ξ_A^ℓ
Type-I	$\cot \beta$	$\cot \beta$	$\cot \beta$	$\cot \beta$	$-\cot \beta$	$-\cot \beta$
Type-II	$\cot \beta$	$-\tan \beta$	$-\tan \beta$	$\cot \beta$	$\tan \beta$	$\tan \beta$
Type-L	$\cot \beta$	$\cot \beta$	$-\tan \beta$	$\cot \beta$	$-\cot \beta$	$\tan \beta$
Type-F	$\cot \beta$	$-\tan \beta$	$\cot \beta$	$\cot \beta$	$\tan \beta$	$-\cot \beta$

Small couplings

- Type-I: easy at large $\tan \beta$ for both A and H
- A: impossible for other 3 types
- H: hard for other 3 types (fine-tuned)

Constraint and 2HDM

Benchmark Scenario:

theoretical constraints, Z-pole direct search, invisible h decay

$$\text{Light } H : \cos(\beta - \alpha) = \frac{1}{\tan \beta}, \quad m_A = m_{H^\pm} = 600 \text{ GeV}, \quad \lambda v^2 = 0 \text{ GeV}^2,$$

$$\text{Light } A : \cos(\beta - \alpha) = 0, \quad m_H = m_{H^\pm} = 90 \text{ GeV}, \quad \lambda v^2 = 0 \text{ GeV}^2 .$$

$$\xi_A^f |_{\cos(\beta-\alpha)=0} = 1/\tan \beta,$$

$$\xi_H^V = c_{\beta-\alpha} = 1/\tan \beta,$$



$$\xi_H^f = c_{\beta-\alpha}(1 - s_{\beta-\alpha}) \approx 1/(2 \tan^3 \beta) + \mathcal{O}(c_{\beta-\alpha}^5)$$

Light H is easier to be long-lived



Results: CP even

$$\xi_H^V = c_{\beta-\alpha} = 1/\tan\beta,$$

$$\xi_H^f = c_{\beta-\alpha}(1 - s_{\beta-\alpha}) \approx 1/(2\tan^3\beta) + \mathcal{O}(c_{\beta-\alpha}^5)$$





Results: CP even



Results: CP odd

$$\xi_A^f |_{\cos(\beta-\alpha)=0} = 1 / \tan \beta$$





Results: for case study

higher luminosity helps to reach the weaker coupling region.

A larger detector, especially the radius helps to extend the reach in mA.


Summary

- General study
 - Production
 - Decay : public code
 - Constraints
- Case study: 2HDM results



Thanks !

Constraint



$$\cos(\beta - \alpha) = 0,$$

$$m_\Phi \equiv m_H = m_A = m_{H^\pm}$$

Theoretical constraints

$$\lambda v^2 \equiv m_\Phi^2 - m_{12}^2 / s_\beta c_\beta$$

$$-125^2 \text{ GeV}^2 < \lambda v^2 < 600^2 \text{ GeV}^2$$



$$\lambda \in (-0.26, 5.95)$$

$$\lambda_4 = \lambda_5 = \lambda_3 - 0.258 = -\lambda$$

Constraint

Theoretical constraints

$$\lambda v^2 \equiv m_H^2 - m_{12}^2/s_\beta c_\beta = 0$$






$$m_H \sim 0 : m_{A/H^\pm} \lesssim 600 \text{ GeV}$$

$$m_A \sim 0 : m_{H^\pm} \lesssim 600 \text{ GeV}, \quad m_H \lesssim m_h$$

Constraint

Oblique constraints: Z pole

Direct search at LEP



$$m_H \sim 0 :$$

$$m_A \sim m_{H^\pm} \lesssim 600 \text{ GeV}$$

$$m_A \sim 0 :$$

$$m_{H^\pm} \sim m_H \lesssim m_h$$

$$m_A \sim 10 \text{ GeV} :$$

$$72 \text{ GeV} \lesssim m_{H^\pm} \sim m_H \lesssim 116 \text{ GeV}$$

Constraint

Invisible Higgs decays

$$\text{Br}(h \rightarrow \phi\phi) = \frac{\Gamma(h \rightarrow \phi\phi)}{\Gamma_h} \approx \frac{1}{\Gamma_h^{\text{SM}}} \frac{g_{h\phi\phi}^2}{8\pi m_h^2} \left(1 - \frac{4m_H^2}{m_h^2}\right)^{1/2} \simeq 4700 \cdot \left(\frac{g_{h\phi\phi}}{v}\right)^2 < 0.24$$



$\text{Br}(h \rightarrow \phi\phi) = 0$

$$\text{Light } H : \cos(\beta - \alpha) = \tan 2\beta \frac{2\lambda v^2 + m_h^2}{2(m_H^2 - 3\lambda v^2 - m_h^2)} \approx \frac{1}{\tan \beta},$$

$$\text{Light } A : \cos(\beta - \alpha) = \tan 2\beta \frac{2\lambda v^2 + m_h^2 + 2m_A^2 - 2m_H^2}{2(m_H^2 - \lambda v^2 - m_h^2)} \approx \frac{1}{\tan \beta} \frac{2m_H^2 - m_h^2}{m_H^2 - m_h^2},$$

Results: CP odd

$$\xi_A^f |_{\cos(\beta-\alpha)=0} = 1/\tan\beta$$



Decay: CP even scalar

$$m_\phi < 2 \text{ GeV}$$

$$H \rightarrow \pi\pi$$

$$H \rightarrow KK$$

$$H \rightarrow \pi\pi\pi\pi$$

$$\begin{aligned} \mathcal{L} &\supset \frac{\Phi}{v} \left(\xi_\Phi^g \frac{\alpha_s}{12\pi} G_{\mu\nu}^a G^{a\mu\nu} - \xi_\Phi^u m_u \bar{u}u - \xi_\Phi^d m_d \bar{d}d - \xi_\Phi^s m_s \bar{s}s \right) \\ &= -\frac{\Phi}{v} \left\{ \xi_\Phi^g \left[\frac{2}{27} \Theta_\mu^\mu - \frac{2}{27} (m_u \bar{u}u + m_d \bar{d}d + m_s \bar{s}s) \right] + (\xi_\Phi^u m_u \bar{u}u + \xi_\Phi^d m_d \bar{d}d + \xi_\Phi^s m_s \bar{s}s) \right\} \end{aligned}$$

$$\Theta_\mu^\mu = -\frac{9\alpha_s}{8\pi} G_{\mu\nu}^a G^{a\mu\nu} + m_u \bar{u}u + m_d \bar{d}d + m_s \bar{s}s.$$

$$\Gamma_\pi = \langle \pi\pi | m_u \bar{u}u + m_d \bar{d}d | 0 \rangle, \quad \Delta_\pi = \langle \pi\pi | m_s \bar{s}s | 0 \rangle, \quad \Theta_\pi = \langle \pi\pi | \Theta_\mu^\mu | 0 \rangle$$


$$\Gamma_{\pi\pi} = \frac{3G_F}{16\sqrt{2}\pi m_\Phi} \beta_\pi \left| \xi_\Phi^{gg} \frac{2}{27} (\Theta_\pi - \Gamma_\pi - \Delta_\pi) + \frac{m_u \xi_\Phi^u + m_d \xi_\Phi^d}{m_u + m_d} \Gamma_\pi + (\xi_\Phi^s) \Delta_\pi \right|^2$$

$$\Gamma_{KK} = \frac{G_F}{4\sqrt{2}\pi m_\Phi} \beta_K \left| \xi_\Phi^{gg} \frac{2}{27} (\Theta_K - \Gamma_K - \Delta_K) + \frac{m_u \xi_\Phi^u + m_d \xi_\Phi^d}{m_u + m_d} \Gamma_K + (\xi_\Phi^s) \Delta_K \right|^2$$


code

- Light Scalar decay
- https://github.com/shiggs90/Light_scalar_decay.git

FASER: Detector



$pp \rightarrow \text{LLP} + X$, LLP travels ~ 480 m, $\text{LLP} \rightarrow \text{charged tracks} + X$



FASER: radius $R = 10$ cm, length $D = 1.5$ m, luminosity $L = 150 \text{ fb}^{-1}$,
FASER 2: radius $R = 1$ m, length $D = 5$ m, luminosity $L = 3 \text{ ab}^{-1}$.

