

Recent Dark Matter combination summary from ATLAS

Ref: <https://arxiv.org/abs/2306.00641>

Results from a wide range of searches targeting different experimental signatures with and without missing transverse momentum (E_T^{miss}) are used to constrain a Two-Higgs-Doublet Model (2HDM) with an additional pseudo-scalar mediating the interaction between ordinary and dark matter (2HDM+a). The analyses use up to 139 fb^{-1} of proton-proton collision data at a centre-of-mass energy $\sqrt{s} = 13\text{ TeV}$ recorded with the ATLAS detector at the Large Hadron Collider between 2015-2018. The results from three of the most sensitive searches are combined statistically. These searches target signatures with large E_T^{miss} and a leptonically decaying Z boson; large E_T^{miss} and a Higgs boson decaying to bottom quarks; and production of charged Higgs bosons in final states with top and bottom quarks, respectively. Constraints are derived for several common as well as new benchmark scenarios within the 2HDM+a.

Primary author(s) : LI, Shu (Tsung-Dao Lee Institute & Shanghai Jiao Tong University); Dr VU, Ngoc Khanh (Tsung-Dao Lee Institute, Shanghai Jiao Tong University)

Presenter(s) : Dr VU, Ngoc Khanh (Tsung-Dao Lee Institute, Shanghai Jiao Tong University)