

Laser-assisted search for axion-like particle or dark photon in strong-field QED

Tong Li (李佟)

Nankai University

based on PRD 111, 055001 (2025), JHEP 07 (2025) 028
with Kai Ma (Xi'an U. Architech. Technol.)


Axion 2025, Nanjing, 2025.7.27-31

Outline

- Strong-field QED and laser-assisted Compton scattering
- Laser-assisted search for axion-like particle and dark photon
- Summary

Strong-field QED

- The quantum field theory in an external and **intense electromagnetic field** is regarded as strong-field QED
- An appropriate theory to study high-intensity physics (unlike high-precision domain, e.g. proton decay)



强场量子真空：施温格极限

- In 1951, J. Schwinger showed that at field strengths of $E = \frac{m_e^2}{e} \sim 1.32 \times 10^{18} \text{ V/m}$, the QED vacuum becomes unstable and decays into electron-positron pairs

$$eE_{cr}\lambda_c = mc^2$$

$$\lambda_c = \hbar/mc = 3.8616 \times 10^{-13} \text{ m} \text{ 电子康普顿波长}$$

正负电子对产生的**施温格极限场强**:

$$E_{cr} = \frac{m^2 c^3}{e\hbar} = 1.323 \times 10^{18} \text{ V/m}$$

对应的**激光强度**:

$$I_{QED} = 2.1 \times 10^{29} \text{ W/cm}^2$$



Phys. Rev. 82 (1951) 664-679

- The calculation of vacuum decay probability exhibits **non-perturbative QED**


$$2VT \frac{(eE)^2}{(2\pi)^3} \sum_{n=1}^{\infty} \frac{1}{n^2} \exp \left[-n\pi \frac{m^2}{eE} \right]$$

- Significant progress has been made in understanding the theory and phenomenology of the Schwinger effect in more realistic backgrounds
- In 1990s, the experiment performed at SLAC observed two strong-field processes in the interaction of an ultra-relativistic electron beam with a terawatt laser pulse


Phys. Rev. Lett. 79, 1626 (1997)

Phys. Rev. D 60, 092004 (1999)


➤ the nonlinear Compton scattering

$$e^{\pm} + n\gamma_L \rightarrow e^{\pm} + \gamma$$


➤ the nonlinear Breit-Wheeler production


$$\gamma + n\gamma_L \rightarrow e^+ + e^-$$


全球10-100PW级超强激光科学装置



credit: Liangliang Ji

全球10-100PW级超强激光科学装置



- Now, the intense electromagnetic field has a lot of applications in atomic physics, nuclear physics and strong-field particle physics
- Two domains:
 1. further studies on nonlinear QED
 2. investigating processes that normally do not occur in vacuum but can be induced under strong fields

Examples:

- Neutron stars strongly magnetized
- Strong magnetic field in high-energy nuclear collisions (e.g. Au-Au at RHIC or Pb-Pb at LHC), CME etc.
- Transmutation of protons in a laser field
 $p \rightarrow n + e^+ + \nu_e$ [New J. Phys. 23, 065007 \(2021\)](#)
- The first laser excitation of the Th-229 low-energy nuclear transition [Phys. Rev. Lett. 132, 182501 \(2024\)](#)
- vacuum birefringence, e.g. Shanghai XFEL
- Search for new physics beyond the SM (this talk)

Laser-induced Compton scattering

- In the presence of an electromagnetic potential, the Dirac equation of a relativistic fermion yields

$$(i\cancel{\partial} - QeA - m)\psi(x) = 0$$

- With circular polarization, the vector potential is

$$A^\mu(\phi) = a_1^\mu \cos(\phi) + a_2^\mu \sin(\phi), \quad \phi = k \cdot x$$

$$a_1^\mu = |\vec{a}|(0, 1, 0, 0), \quad a_2^\mu = |\vec{a}|(0, 0, 1, 0)$$

$$a = \varepsilon_0 / \omega$$

the electric field strength the laser frequency

- The wave function of electron is given by the Volkov state
- dressed state by the interaction between electron and laser photons

$$\psi_{p,s}(x) = \left[1 + \frac{Qe \not{k} A}{2(k \cdot p)} \right] \frac{u(p, s)}{\sqrt{2q^0 V}} e^{iF_1(q,s)} \quad \text{Z. Physik 94, 250 (1935)}$$

$$F_1(q, s) = -q \cdot x - \frac{Qe(a_1 \cdot p)}{(k \cdot p)} \sin \phi + \frac{Qe(a_2 \cdot p)}{(k \cdot p)} \cos \phi$$

- effective momentum and mass of the dressed electron

$$q^\mu = p^\mu + \frac{Q^2 e^2 a^2}{2k \cdot p} k^\mu$$

$$q^2 = m_e^2 + Q^2 e^2 a^2 = m_e^{*2}$$

- High-order (nonlinear) effects in the laser field are included

- Consider the laser-induced Compton scattering

$$e^-(p) + n\omega(k) \rightarrow e^-(p') + \gamma(k')$$

- The S matrix

$$S_{fi} = ie \frac{1}{\sqrt{2k'^0 V}} \int d^4x e^{ik' \cdot x} \overline{\psi_{p',s'}(x)} \not{\epsilon} \psi_{p,s}(x)$$

$$\begin{aligned} \mathcal{M} &= ie \frac{1}{\sqrt{2k'^0 V}} e^{ik' \cdot x} \overline{\psi_{p',s'}(x)} \not{\epsilon} \psi_{p,s}(x) \\ &= ie \frac{e^{i(k'+q'-q) \cdot x} \boxed{e^{-i\Phi}}}{\sqrt{2^3 V^3 q^0 q'^0 k'^0}} \overline{u(p', s')} \left[1 - \frac{e \not{A} \not{k}}{2k \cdot p'} \right] \not{\epsilon} \left[1 - \frac{e \not{k} \not{A}}{2k \cdot p} \right] u(p, s) \end{aligned}$$

$$\Phi = ea_1 \cdot y \sin \phi - ea_2 \cdot y \cos \phi \qquad y^\mu = \frac{p'^\mu}{k \cdot p'} - \frac{p^\mu}{k \cdot p}$$

- The amplitude can be written as

$$\mathcal{M} = ie \frac{e^{i(k'+q'-q)\cdot x}}{\sqrt{2^3 V^3 q^0 q'^0 k'^0}} \sum_{i=0}^2 C_i \mathcal{M}_i$$

$$\begin{aligned} \mathcal{M}_0 &= \overline{u}_2 \not{\epsilon} u_1 + \overline{u}_2 \frac{e^2 a^2}{2k \cdot p k \cdot p'} k \cdot \epsilon \not{k} u_1 & C_0 &= e^{-i\Phi} , \\ \mathcal{M}_1 &= -\overline{u}_2 \not{\epsilon} \frac{e \not{k} \not{a}_1}{2k \cdot p} u_1 - \overline{u}_2 \frac{e \not{a}_1 \not{k}}{2k \cdot p'} \not{\epsilon} u_1 & C_1 &= \cos \phi e^{-i\Phi} \\ \mathcal{M}_2 &= -\overline{u}_2 \not{\epsilon} \frac{e \not{k} \not{a}_2}{2k \cdot p} u_1 - \overline{u}_2 \frac{e \not{a}_2 \not{k}}{2k \cdot p'} \not{\epsilon} u_1 & C_2 &= \sin \phi e^{-i\Phi} \end{aligned}$$

- Look into the phase function $\Phi = ea_1 \cdot y \sin \phi - ea_2 \cdot y \cos \phi$

$$\Phi = z \sin(\phi - \phi_0)$$

$$z = e \sqrt{(a_1 \cdot y)^2 + (a_2 \cdot y)^2}, \quad \cos \phi_0 = \frac{ea_1 \cdot y}{z}, \quad \sin \phi_0 = \frac{ea_2 \cdot y}{z}$$

- Then

$$e^{-i\Phi} = e^{-iz \sin(\phi - \phi_0)} = \sum_{n=-\infty}^{\infty} c_n e^{-in(\phi - \phi_0)}$$

$$c_n = \frac{1}{2\pi} \int_{-\pi}^{\pi} d\varphi e^{-iz \sin \varphi} e^{in\varphi} = J_n(z)$$

- Finally (Jacobi-Anger expansion)

$$C_0 = e^{-i\Phi} = \sum_{n=-\infty}^{\infty} B_n(z) e^{-in\phi} \quad B_n(z) = J_n(z) e^{in\phi_0}$$


$$\cos \phi = \frac{1}{2} \left(e^{i\phi} + e^{-i\phi} \right), \quad \sin \phi = \frac{1}{2i} \left(e^{i\phi} - e^{-i\phi} \right)$$

$$C_1 = \cos \phi e^{-i\Phi} = \frac{1}{2} \sum_{n=-\infty}^{\infty} \left[B_{n+1}(z) + B_{n-1}(z) \right] e^{-in\phi}$$

$$C_2 = \sin \phi e^{-i\Phi} = \frac{1}{2i} \sum_{n=-\infty}^{\infty} \left[B_{n+1}(z) - B_{n-1}(z) \right] e^{-in\phi}$$

- Combine $e^{-in\phi}$ with other plane waves

$$\mathcal{M} = ie \sum_{n=-\infty}^{\infty} \frac{e^{i(k'+q'-q-nk)\cdot x}}{\sqrt{2^3 V^3 q^0 q'^0 k'^0}} \sum_{i=0}^2 \tilde{C}_i^n \mathcal{M}_i$$

B functions 

- Integrating out the coordinate x

$$S_{fi} = ie \sum_{n=-\infty}^{\infty} \frac{(2\pi)^4 \delta^4(\overbrace{k' + q'}^{\text{final}} - \overbrace{q + nk}^{\text{initial}})}{\sqrt{2^3 V^3 q^0 q'^0 k'^0}} \sum_{i=0}^2 \tilde{C}_i^n \mathcal{M}_i$$

- Squared S matrix and scattering cross section


$$|S_{fi}|^2 = e^2 \sum_{n=-\infty}^{\infty} \frac{(2\pi)^4 \delta^4(k' + q' - q - nk) VT}{2^3 V^3 q^0 q'^0 k'^0} \sum_{i,j=0}^2 \tilde{C}_i^n (\tilde{C}_j^n)^\dagger \overline{\mathcal{M}_i \mathcal{M}_j^\dagger}$$

$$\begin{aligned} \sigma &= \frac{|S_{fi}|^2}{VT} \frac{1}{2(1/V)} \frac{1}{\rho_\omega} V \int \frac{d^3 q'}{(2\pi)^3} V \int \frac{d^3 k'}{(2\pi)^3} \\ &= \frac{1}{2\rho_\omega} \frac{e^2}{2q^0} \sum_{n=-\infty}^{\infty} \int d\Pi_2 \sum_{i,j=0}^2 \tilde{C}_i^n (\tilde{C}_j^n)^\dagger \overline{\mathcal{M}_i \mathcal{M}_j^\dagger} \end{aligned}$$

$$\sum_{i,j=0}^2 \tilde{C}_i^n (\tilde{C}_j^n)^* \overline{\mathcal{M}_i \mathcal{M}_j^\dagger} = -4m_e^2 J_n^2(z) + e^2 a^2 \frac{1+u^2}{u} [J_{n-1}^2(z) + J_{n+1}^2(z) - 2J_n^2(z)]$$

$$u \equiv k \cdot p / k \cdot p' \quad z = \frac{2nuea}{s'_n - m_e^{*2}} \left(\frac{s'_n + m_e^{*2} - m_\chi^2}{u} - \frac{s'_n}{u^2} - m_e^{*2} \right)^{1/2}$$

- define an intensity quantity $\eta \equiv \frac{ea}{m_e} = \frac{e\mathcal{E}_0}{\omega_{\text{Lab}} m_e}$
- power series expansion of η
- This result is nonperturbative in character and the nonlinear effects become important when $\eta \gtrsim 1$
- The cross section is in unit of barn!



Laser-assisted search for dark particles

- dark photon (ALP): $U(1)_D$

$$\mathcal{L}_{\text{DP}} \supset -\frac{1}{4}F^{\mu\nu}F_{\mu\nu} - \frac{1}{4}F_D^{\mu\nu}F_{D\mu\nu} - \frac{\epsilon}{2}F^{\mu\nu}F_{D\mu\nu} + \frac{1}{2}m_D^2 A_D^\mu A_{D\mu} \\ - e\epsilon J_{\text{EM}}^\mu A_{D\mu} = -eQ\epsilon \bar{\psi}\gamma^\mu\psi A_{D\mu}$$

- axion-like particle (ALP): pseudo-NG boson

$$\mathcal{L}_{\text{ALP}} \supset c_{ae} \frac{\partial_\mu a}{2f_a} \bar{e}\gamma^\mu\gamma_5 e$$

$$g_{ae} = c_{ae} m_e / f_a$$

- hypothesis: invisible for m_a or $m_{\gamma_D} < 1 \text{ MeV}$

- Consider the **Compton scattering to DP or ALP**

$$e^-(p) + n\omega(k) \rightarrow e^-(p') + \boxed{\gamma_D/a(k')}$$

- The new amplitude square

DP:
$$\sum_{i,j=0}^2 \tilde{C}_i^m (\tilde{C}_j^m)^* \overline{\mathcal{M}_i^{\gamma_D} \mathcal{M}_j^{\gamma_D \dagger}} = -2(2m_e^2 + m_\chi^2) J_n^2(z) \\ + e^2 a^2 \frac{1+u^2}{u} [J_{n-1}^2(z) + J_{n+1}^2(z) - 2J_n^2(z)]$$


ALP:
$$\sum_{i,j=0}^2 \tilde{C}_i^m (\tilde{C}_j^m)^* \overline{\mathcal{M}_i^a \mathcal{M}_j^{a \dagger}} = -4m_e^2 m_\chi^2 J_n^2(z) \\ + e^2 a^2 \frac{2m_e^2 (1-u)^2}{u} [J_{n-1}^2(z) + J_{n+1}^2(z) - 2J_n^2(z)]$$

- Experimental setup (e.g. SLAC)

$$E_{Lab} = 46.6 \text{ GeV}$$



$$\omega_{Lab} = 2.35 \text{ eV}$$

$$\theta_{Lab} = 17^\circ$$





- Cross section

Kai Ma, TL, PRD 111, 055001 (2025)



- Distribution of outgoing electron energy: edges





Kai Ma, TL, PRD 111, 055001 (2025)

- Sensitivity of laser-induced Compton scattering to dark particle couplings (SM bkg: $e^- + \text{laser} \rightarrow e^- + \nu + \bar{\nu}$)

$$\frac{S}{\sqrt{S+B}}$$

- Complementary to other beam dump and collider experiments

Kai Ma, TL, PRD 111, 055001 (2025)



Laser-induced Compton scattering to DM in EFT

Kai Ma, TL, JHEP 07 (2025) 028

- Dirac-type fermionic DM in a leptophilic scenario

$$e^- + \text{laser} \rightarrow e^- + \chi + \bar{\chi}$$

$$\mathcal{O}_{MD} = (\bar{\chi} \sigma^{\mu\nu} \chi) F_{\mu\nu} , \quad \mathcal{O}_{ED} = (\bar{\chi} \sigma^{\mu\nu} i \gamma_5 \chi) F_{\mu\nu} ,$$

$$\mathcal{O}_{SS} = (\bar{e} e) (\bar{\chi} \chi) , \quad \mathcal{O}_{SP} = (\bar{e} e) (\bar{\chi} i \gamma_5 \chi) ,$$



$$\mathcal{O}_{PS} = (\bar{e} i \gamma_5 e) (\bar{\chi} \chi) , \quad \mathcal{O}_{PP} = (\bar{e} i \gamma_5 e) (\bar{\chi} i \gamma_5 \chi) ,$$

$$\mathcal{O}_{VV} = (\bar{e} \gamma^\mu e) (\bar{\chi} \gamma_\mu \chi) , \quad \mathcal{O}_{VA} = (\bar{e} \gamma^\mu e) (\bar{\chi} \gamma_\mu \gamma_5 \chi) ,$$

$$\mathcal{O}_{AV} = (\bar{e} \gamma^\mu \gamma_5 e) (\bar{\chi} \gamma_\mu \chi) , \quad \mathcal{O}_{AA} = (\bar{e} \gamma^\mu \gamma_5 e) (\bar{\chi} \gamma_\mu \gamma_5 \chi) .$$

- Sensitivity of laser-induced Compton scattering to the effective cutoff scale

Kai Ma, TL, JHEP 07 (2025) 028



Summary

- The laser of an intense electromagnetic field plays as an important tool to study the strong-field particle physics and search for new physics beyond the SM
- We investigate the laser-induced Compton scattering to dark particles such as invisible dark photon or axion-like particle
- We find that the laser-induced process provides a complementary and competitive search of new dark particles lighter than 1 MeV

Summary

- The laser of an intense electromagnetic field plays as an important tool to study the strong-field particle physics and search for new physics beyond the SM
- We investigate the laser-induced Compton scattering to dark particles such as invisible dark photon or axion-like particle
- We find that the laser-induced process provides a complementary and competitive search of new dark particles lighter than 1 MeV

Thank you!