

# What do experimental data tell us?

Junjie Cao

junjiec@alumni.itp.ac.cn

Henan Normal University,  
Department of Physics

August 2, 2022

**Conjectures on the natural realization of WIMP DM  
in economic SUSY; Studies initiated in 2016.**

DM-EFTs	Example	Annihilation	$\tilde{\chi} - N$ Scattering	Remarks
SM+DM	SM+ $S_{real}$	Weak/contact interactions	$\sigma_{SI} \gtrsim 10^{-45} \text{cm}^2$ and/or $\sigma_{SD} \gtrsim 10^{-39} \text{cm}^2$	Experimentally excluded.
			Suppressed by cancellation	Symmetry!
		Feeble interaction: $h/Z$ funnels	Suppressed	Fine-tuning: $\Delta > 100$ .
SM+DM+X	MSSM with Light Gauginos	Coannihilation	Suppressed	Fine-tuning: $\Delta > 30$ ; Tight LHC constraints.
SM+DM+XY	GNMSSM ISS-NMSSM	May form <b>secluded DM sector</b>	Suppressed	No tuning; <b>three portals to SM.</b>

## WIMP DM:

**Weak interactions in the DM sector and feeble interactions between SM and DM sectors.**

## Motivations



- Natural EW symmetry breaking: **MSSM**  $\rightarrow$   **$Z_3$ -NMSSM**  $\rightarrow$  **General NMSSM**.
- Neutrino mass: **Type-I NMSSM**  $\rightarrow$  **ISS-NMSSM**  $\rightarrow$  **B-L NMSSM**.

- 1 Preliminaries of WIMP DM (excerpt from Jia Liu's talk)
- 2 Example I: MSSM
- 3 Example II:  $Z_3$ -NMSSM
- 4 Section III: General NMSSM
- 5 Section IV: Type-I NMSSM
- 6 Example V: B-L NMSSM
- 7 Conclusions

# Section I

## Preliminaries of WIMP DM

# Preliminary: the freeze-out of thermal DM





- Mass bound:  $N_{\text{eff}}$  from CMB, unitary  
 $5 \text{ MeV} \lesssim m_{\text{DM}} \lesssim 110 \text{ TeV}$ ;
- DM starts with thermal distribution;
- Relic abundance is determined by freeze-out mechanism;
- DM has an electroweak-scale coupling (WIMP miracle).

Consider  $\text{DM DM} \rightarrow \text{X X}$ :

$$\langle \sigma v \rangle \sim \frac{g^4}{m_{\text{DM}}^2} \sim 3 \times 10^{-26} \text{ cm}^3 \text{ s}^{-1} \Rightarrow g \sim \sqrt{\frac{m_{\text{DM}}}{10 \text{ TeV}}},$$

$$g \sim 0.1 \text{ for } m_{\text{DM}} = 100 \text{ GeV}.$$

# Preliminary: limits from direct detection experiments



arXiv:1805.12562


arXiv:1902.03234

**Preliminary PandaX-4T results released!**


# Preliminary: The way-out from direct detection limits

- 1. Very small coupling:

- 1.1 Secluded dark matter (dark sector)



Proposed in 0711.4866.  
Three types of portals:  
Higgs portal;  
Gauge portal;  
Neutrino portal



**Dark mediator  
with very small coupling to SM**

**This mechanism can be realized in non-minimal SUSY!**

# Preliminary: The way-out from direct detection limits

## • 2. Suppressed scattering cross-section:

### • By velocity or momentum transfer

Case for Fermionic DM

Kumar & Marfatia:1305.1611 (PRD)


	Name	Interaction Structure	$\sigma_{SI}$ suppression	$\sigma_{SD}$ suppression	$s$ -wave?
Scalar	F1	$\bar{X} X \bar{q} q$	1	$q^2 v^{\perp 2}$ (SM)	No
	F2	$\bar{X} \gamma^5 X \bar{q} q$	$q^2$ (DM)	$q^2 v^{\perp 2}$ (SM); $q^2$ (DM)	Yes
	F3	$\bar{X} X \bar{q} \gamma^5 q$	0	$q^2$ (SM)	No
Pseudoscalar	F4	$\bar{X} \gamma^5 X \bar{q} \gamma^5 q$	0	$q^2$ (SM); $q^2$ (DM)	Yes
Vector	F5	$\bar{X} \gamma^\mu X \bar{q} \gamma_\mu q$ (vanishes for Majorana $X$ )	1	$q^2 v^{\perp 2}$ (SM) $q^2$ (SM); $q^2$ or $v^{\perp 2}$ (DM)	Yes
	F6	$\bar{X} \gamma^\mu \gamma^5 X \bar{q} \gamma_\mu q$	$v^{\perp 2}$ (SM or DM)	$q^2$ (SM)	No
Anapole	F7	$\bar{X} \gamma^\mu X \bar{q} \gamma_\mu \gamma^5 q$ (vanishes for Majorana $X$ )	$q^2 v^{\perp 2}$ (SM); $q^2$ (DM)	$v^{\perp 2}$ (SM) $v^{\perp 2}$ or $q^2$ (DM)	Yes
	F8	$\bar{X} \gamma^\mu \gamma^5 X \bar{q} \gamma_\mu \gamma^5 q$	$q^2 v^{\perp 2}$ (SM)	1	$\propto m_t^2/m_X^2$
	F9	$\bar{X} \sigma^{\mu\nu} X \bar{q} \sigma_{\mu\nu} q$ (vanishes for Majorana $X$ )	$q^2$ (SM); $q^2$ or $v^{\perp 2}$ (DM) $q^2 v^{\perp 2}$ (SM)	1	Yes
	F10	$\bar{X} \sigma^{\mu\nu} \gamma^5 X \bar{q} \sigma_{\mu\nu} q$ (vanishes for Majorana $X$ )	$q^2$ (SM)	$v^{\perp 2}$ (SM) $q^2$ or $v^{\perp 2}$ (DM)	Yes

**Not easy to build specific models! Let alone in supersymmetric**




# Preliminary: The way-out from direct detection limits

## • 3. Coannihilation mechanism



- $\tilde{Y}$  has a close mass with DM
- $\tilde{Y}$  is not populated today due to decay
- Charged  $\tilde{Y}$ : near degenerate spectrum of SUSY, AMSB; EW multiplet DM  $(2n+1, 0)$  ( $\delta m \sim 166$  MeV)



Easily realized in SUSY, but unnatural!



# Preliminary: The way-out from direct detection limits

## • 4. Resonant annihilation

- $2m_{\text{DM}} \approx m_X$

Scalar DM ( $s$ ) with a Higgs portal coupling

$$\Delta\mathcal{L}_s = -\frac{1}{2}m_s^2 s^2 - \frac{1}{4}\lambda_s s^4 - \frac{1}{4}\lambda_{Hss}\phi^\dagger\phi s^2$$



Arcadi et al: 2101.02507

See also WL Guo, LY Wu et al 2010; B Li, YF Zhou 2015


**Easily realized in SUSY, but needs severe fine-tuning!**

# Preliminary: The way-out from direct detection limits

## • 5. Cancellation effect in scattering cross-section

### • SM Higgs - Dark scalar mediator cancellation

Gross, Lebedev1, Toma: 1708.02253 (PRL)



See JL, XP Wang and F Yu 1704.00730 (JHEP),  
for cancellation between A' - Z boson in kinetic  
mixing dark photon model

$$V_0 = -\frac{\mu_H^2}{2} |H|^2 - \frac{\mu_S^2}{2} |S|^2 + \frac{\lambda_H}{2} |H|^4 + \lambda_{HS} |H|^2 |S|^2 + \frac{\lambda_S}{2} |S|^4$$

$$V_{\text{soft}} = -\frac{\mu_S^2}{4} S^2 + \text{h.c.} \quad \text{symmetry : } S \leftrightarrow S^*$$

$$S = (v_s + s + i\chi)/\sqrt{2} \quad \text{Pseudoscalar DM}$$

CP-even scalar mixing (s, h)  $\rightarrow$  ( $h_1, h_2$ )

$$\mathcal{L} \supset -(h_1 \cos \theta + h_2 \sin \theta) \sum_f \frac{m_f}{v} \bar{f} f \quad \mathcal{L} \supset \frac{\kappa^2}{2v_s} \left( m_{h_1}^2 \sin \theta h_1 - m_{h_2}^2 \cos \theta h_2 \right)$$

$$A_{dd}(t) \propto \sin \theta \cos \theta \left( \frac{m_{h_2}^2}{t - m_{h_2}^2} - \frac{m_{h_1}^2}{t - m_{h_1}^2} \right) \simeq \sin \theta \cos \theta \frac{t(m_{h_2}^2 - m_{h_1}^2)}{m_{h_1}^2 m_{h_2}^2} \simeq 0$$


The amplitude is suppressed by  $q^2$

**Can not be realized in SUSY!**

# Preliminary: The way-out from direct detection limits

## • 6. Leptophilic models

- Only couples to electrons, couples to nucleons at 1-loop
- For light DM, e-DM recoils can have stringent limits (e.g. XENON1T, PANDAX, CDEX)
- For heavy DM, nucleus-DM recoils wins over e-DM recoil



$$R^{\text{WAS}} : R^{\text{WES}} : R^{\text{WNS}} \sim \epsilon_{\text{WAS}} : \epsilon_{\text{WES}} \frac{m_e}{m_N} : \left( \frac{\alpha_{\text{em}} Z}{\pi} \right)^2 \sim 10^{-17} : 10^{-10} : 1$$

WAS = e kicked out

WES = e to higher energy level

WNS = nucleus recoil

The probability to find a high p electron in the wave function is highly suppressed!

Kopp et al: 0907.3159 (PRD)

Realized the SM extensions with  $L_{\mu-\tau}$ ,  $B - L$ , or left-right symmetry, and their supersymmetric versions!

# Preliminary: indirect detection from DM annihilation

- observable quantity:

CMB photon, photon from dwarf galaxies, and positron from cosmic ray, etc.

- $S \propto \sum_i \langle \sigma v \rangle_{0,i} \epsilon_i,$

$\langle \sigma v \rangle_{0,i}$ : annihilation rate for  $\text{DM DM} \rightarrow e^+e^-, \mu^+\mu^-, \tau^+\tau^-, t\bar{t}, \dots$   
at present day;

$\epsilon_i$ : efficiency translating annihilation products into signal.

Note:  $\langle \sigma v \rangle_F$  may differ significantly from  $\langle \sigma v \rangle_{0,i}$

$$\sigma v \sim \sigma_s + \sigma_p v^2 + \sigma_d v^4 + \dots \text{ (s-, p-, and d-wave contribution)}$$

- Freeze-out:  $v^2 \sim 0.25$
- CMB:  $v^2 \sim \text{eV}/m_{\text{DM}} \sim 10^{-5}$
- Today:  $v \sim 10^{-3}c$

$\epsilon_i$  may differ greatly for different annihilation final state!

# Preliminary: indirect detection from DM annihilation




Figure 1: Planck CMB limits at 95% C.L. for DM annihilation 100% to individual channels.





Figure 2: Fermi-LAT limits at 95% C.L. for DM annihilation 100% to individual channels.

**Depending on final state, CMB limits are powerful for light DM, while Fermi-LAT limits are effective for  $m_{\text{DM}} \lesssim 100$  GeV.**

# Preliminary: indirect detection from DM annihilation



**Figure 3:** Bounds on the generic thermal WIMP window (s-wave  $2 \rightarrow 2$  annihilation, standard cosmological history), assuming WIMP DM is 100% of the DM. Shown is the conservative bound calculated from the data of CMB, Fermi-LAT and AMS-02 (Visibles), and the unitarity bound. The remaining WIMP window is the orange line, and the white space is unprobed. Thermal relic cross section is the dashed line.

# Conclusion about WIMP DM

- ① As far as WIMP DM itself is concerned, it can fit experiments very well.
- ② WIMP DM can be easily embedded into renormalizable theories. In this case, DM physics usually entangles with Higgs physics, sparticle physics, and sometimes neutrino physics. Global fit is necessary.
- ③ In economic WIMP DM theories, DM physics are usually in tension with various experiments, and consequently, the theories become unnatural.
- ④ What is the most economic and natural (supersymmetric) WIMP DM theory?



## Section II

### Criteria in Estimating the Goodness of a Theory

# CEGT: How to study SUSY phenomenology?

Many many ways! Most advanced method:

**Fit theory to experimental data, extract underlying physics!**

- ① Construct **likelihood** function from **experimental data**;
- ② Scan theory's parameter space with **advanced algorithm**:
  - Characteristics of the parameter space:  
high dimensional, highly degenerated likelihood, isolated physical parameter island, inefficient for random and Markov chain scan;
  - **MultiNest algorithm** is well adaptive for such a situation:  
use *nlive* samples to decide iso-likelihood contour in each iteration;  
provide comprehensive information of the space;  
the results are statistically significant. **Why?  $\Rightarrow$**
- ③ Scrutinize the **properties** of obtained parameter points, e.g., prediction on various experimental measurements, theoretical fine-tuning, vacuum stability, etc.. **Human learning materials!**
- ④ Analyze global features of the theory by **statistics**:  
**Some fundamental physical mechanisms can be inferred.**
- ⑤ Provide intuitive **understand**. with the help of analy. formulae.

**Bayesian theorem:**  $\Theta = (\Theta_1, \Theta_2, \dots)$  is theoretical input parameters.

$$P(\Theta | \mathbf{D}, H) \equiv \frac{P(\mathbf{D} | \Theta, H) P(\Theta | H)}{P(\mathbf{D} | H)} \implies P(\Theta) \equiv \frac{\mathcal{L}(\Theta) \pi(\Theta)}{\mathcal{Z}}$$

- $P(\Theta | \mathbf{D}, H) \equiv P(\Theta)$ : Posterior probability distribution function.
- $P(\mathbf{D} | \Theta, H) \equiv \mathcal{L}(\Theta)$ : Likelihood function.
- $P(\Theta | H) \equiv \pi(\Theta)$ : Prior probability Density Function.
- $P(\mathbf{D} | H) \equiv \mathcal{Z}$ : Bayesian evidence, normalization factor.

$P(\Theta)$ : the state of our knowledge about the parameters  $\Theta$  given the experimental data  $D$ , or alternatively speaking, the updated prior PDF after considering the impact of the experimental data.

One can infer from  $P(\Theta)$  the underlying physics of the model.

**Likelihood function:** the preference of experimental results to parameter point  $p = \{\Theta\}$ , e.g., Gaussian distribution:

$$\mathcal{L} = e^{-\frac{[\mathcal{O}_{th}(\Theta) - \mathcal{O}_{exp}]^2}{2\sigma^2}}.$$

$\mathcal{O}_{th}(\Theta)$ : theoretical prediction,       $\mathcal{O}_{exp}$ : experimental measurement,  
 $\sigma$ : total uncertainty.

**Bayesian evidence:** averaged likelihood, reflecting theory's capability to keep consistent with the data.

$$\mathcal{Z} = \int \mathcal{L}(\Theta) \pi(\Theta) d^D \Theta.$$

$D$ : Dimension of the parameter space.

**Marginal posterior PDFs:** reflecting the preference to specific regions of one or more parameters.

$$1D : P(\Theta_A) = \int P(\Theta) d\Theta_1 d\Theta_2 \cdots d\Theta_{A-1} d\Theta_{A+1} \cdots$$

$$2D : P(\Theta_A, \Theta_B) = \int P(\Theta) d\Theta_1 d\Theta_2 \cdots d\Theta_{A-1} d\Theta_{A+1} \cdots d\Theta_{B-1} d\Theta_{B+1} \cdots$$

**Credible Regions:** most preferred parameter regions by data; **it depends on both likelihood function and phase space.**

$$1D : \int_{\Theta_{A_1}}^{\Theta_{A_2}} P(\Theta_A) d\Theta_A = 1 - \alpha$$

$$2D : \int_{P(\Theta_A, \Theta_B) \geq p_{\text{crit}}} P(\Theta_A, \Theta_B) d\Theta_A d\Theta_B = 1 - \alpha$$

$$1\sigma: \alpha = 0.317, \quad 2\sigma: \alpha = 0.055.$$

**Profile Likelihood:** parameter's capability to explain the data.

$$1D : \mathcal{L}(\Theta_A) = \max_{\Theta_1, \dots, \Theta_{A-1}, \Theta_{A+1}, \dots} \mathcal{L}(\Theta),$$

$$2D : \mathcal{L}(\Theta_A, \Theta_B) = \max_{\Theta_1, \dots, \Theta_{A-1}, \Theta_{A+1}, \dots, \Theta_{B-1}, \Theta_{B+1}, \dots} \mathcal{L}(\Theta)$$

**Confidence Intervals:** most favored regions to explain the data; **it depends only on the likelihood function.**

$$1D : \{ \chi^2(\Theta_A) - \chi_{Best}^2 \} \leq F_{\chi_1^2}^{-1}(1 - \alpha),$$

$$2D : \{ \chi^2(\Theta_A, \Theta_B) - \chi_{Best}^2 \} \leq F_{\chi_2^2}^{-1}(1 - \alpha)$$

$$\chi^2(\Theta_A) \equiv -2 \log \mathcal{L}(\Theta_A), \quad \chi^2(\Theta_A, \Theta_B) \equiv -2 \log \mathcal{L}(\Theta_A, \Theta_B);$$

$\chi_{Best}^2$ : the  $\chi^2$  value for the best point;

$F_{\chi_n^2}^{-1}$ : the inverse cdf for a chi-squared distribution with n dof:

$$1\sigma \ (\alpha = 0.317): F_{\chi_1^2}^{-1} = 1.00, \quad F_{\chi_2^2}^{-1} = 2.30;$$



$$2\sigma \ (\alpha = 0.046): F_{\chi_1^2}^{-1} = 4.00, \quad F_{\chi_2^2}^{-1} = 6.18.$$

## Rich experimental data have been accumulated !

- Precision electroweak data;
- Heavy flavor data;
- Neutrino experiments;
- Higgs property measurement.
- **Dark matter search experiments;**
- **LHC search for supersymmetry;**
- **Muon anomalous magnetic moment.**

**Global Fit: Combine all the data to analyze theories.**

# CEGT: WIMP DM direct search experiments



arXiv:1805.12562

arXiv:1902.03234

**Preliminary PandaX-4T results released!**





# CEGT: Implications of DM search experiments

**Popular WIMP DM candidate: Bino-dominated  $\tilde{\chi}_1^0$  in MSSM.**  
DM-nucleon scatterings proceed by *t*-channel exchange of Higgs and *Z* boson, respectively.

$$\sigma_{\tilde{\chi}_1^0-N}^{\text{SI}} \simeq 5 \times 10^{-45} \text{ cm}^2 \left( \frac{C_{\tilde{\chi}_1^0 \tilde{\chi}_1^0 \text{h}}}{0.1} \right)^2 \left( \frac{m_{\text{h}}}{125 \text{ GeV}} \right)^2$$
$$\sigma_{\tilde{\chi}_1^0-N}^{\text{SD}} \simeq 10^{-39} \text{ cm}^2 \left( \frac{C_{\tilde{\chi}_1^0 \tilde{\chi}_1^0 \text{Z}}}{0.1} \right)^2$$

- ❶ Interactions of DM with SM particles are **feeble at most** when  $m_{\tilde{\chi}_1^0} \sim 100 \text{ GeV}$ .
- ❷ Difficult to obtain the measured abundance if  $\text{DM DM} \rightarrow \text{SM SM}$ .  
Exceptions: Co-annihilation, Resonance annihilation.  
**All corresponds to a small Bayesian evidence, fine-tuned!**
- ❸ **Simple WIMP DM theories are becoming unnatural!**
- ❹ Good theory: Naturally explaining the experimental results.  
E.g., secluded DM theories in a more complex framework.


# CEGT: LHC searches for SUSY



## Latest LHC searches for tri- and bi-lepton signals.

- 1 Simplified model for a specified process.
- 2 Invalid for a specific theory: complex decay chain, multiple production processes, and various signals to be analyzed.
- 3 Elaborated Monte Carlo simulations are necessary.

## Muon g – 2 in SUSY:




**Operator contributing to  $a_\mu$ :**  $\frac{a_\mu}{m_\mu} \bar{b} \sigma_{\mu\nu} b F^{\mu\nu}$ .

**Note that the operator involves the chiral flipping of  $\mu$  leptons.**

# CEGT: Improved measurement of Muon g-2

After considering experimental constraints on sparticle masses, mass insertion method is a very good approximation.

$$\Delta a_\mu = a_\mu^{\text{Exp}} - a_\mu^{\text{SM}} = (251 \pm 59) \times 10^{-11}$$



- ① Moderately **large  $\tan \beta$**  is necessary!
- ② Set an **upper bound** on LSP mass (about 600 GeV).
- ③ Set an **upper bound** on NLSP mass (about 700 GeV).
- ④ The other involved sparticles **cannot be excessively heavy**.
- ⑤ LHC search **tightly limits** SUSY explanantions:  $1 + 1 \gg 2$ .
- ⑥ Global fits with/without Muon g-2 **differs** significantly.

Readily explain data, particularly those for correlated obs.!

- ① **DM physics:**  $\Omega h^2$  **versus**  $\sigma_{\tilde{\chi}_1^0-p}$  / Bayesian Evidence.  
 $\times/\checkmark$ : explain the experimental results with/**without** tuning.
- ② **LHC and  $\Delta a_\mu$ :** SUSY searches **versus** sizable correction to  $a_\mu$ .  
 $\times/\checkmark$ : tight/**loose** constraints on the explanation of Muon g-2.
- ③ **Natural EWSB:**  $m_Z^2 = 2(m_{H_d}^2 - m_{H_u}^2 \tan^2 \beta)/(\tan^2 \beta - 1) - 2\mu^2$ .  
 $\times/\checkmark$ : whether or not a moderately **small**  $\mu$  is preferred.
- ④ **Neutrino physics:** neutrino masses and mixings.  
 $\times/\checkmark$ : whether or not **providing** reasonable mechanisms for ....
- ⑤ **Higgs physics:** unreasonably large mass, SM-like couplings  
 $\times/\checkmark$ : explain the mass with/**without** large radiative corrections.

# CEGT: Status of different supersymmetric theories

Obtained by both analytic formulae and global fits, which are tough tasks.

Model	MSSM	$Z_3$ -NMSSM		GNMSSM	Type-I/ISS NMSSM	B-L NMSSM
DM component	Bino	Bino	Singlino	Singlino	Sneutrino	Bino, Singlino, BLino, Bileptino, or sneutrino
DM physics	×	×	×	✓	✓	✓
LHC and $\Delta a_\mu$	×	×	×	✓	✓	✓
EWSB	×	×	✓	✓	✓	✓
Neutrino	×	×	×	×	✓	✓
Higgs mass	×	×	×	×	✓	✓

## Section III

### Example I: MSSM

# MSSM: Theoretical preliminaries

- Vector Superfields

SF	Spin $\frac{1}{2}$	Spin 1	$SU(N)$	Coupling	Name
$\hat{B}$	$\lambda_{\hat{B}}$	$B$	U(1)	$g_1$	hypercharge
$\hat{W}$	$\lambda_{\hat{W}}$	$W$	SU(2)	$g_2$	left
$\hat{g}$	$\lambda_{\hat{g}}$	$g$	SU(3)	$g_3$	color

- Chiral Superfields

SF	Spin 0	Spin $\frac{1}{2}$	Generations	$(U(1) \otimes SU(2) \otimes SU(3))$
$\hat{q}$	$\tilde{q}$	$q$	3	$(\frac{1}{6}, \mathbf{2}, \mathbf{3})$
$\hat{l}$	$\tilde{l}$	$l$	3	$(-\frac{1}{2}, \mathbf{2}, \mathbf{1})$
$\hat{H}_d$	$H_d$	$\tilde{H}_d$	1	$(-\frac{1}{2}, \mathbf{2}, \mathbf{1})$
$\hat{H}_u$	$H_u$	$\tilde{H}_u$	1	$(\frac{1}{2}, \mathbf{2}, \mathbf{1})$
$\hat{d}$	$\tilde{d}_R^*$	$d_R^*$	3	$(\frac{1}{3}, \mathbf{1}, \mathbf{\bar{3}})$
$\hat{u}$	$\tilde{u}_R^*$	$u_R^*$	3	$(-\frac{2}{3}, \mathbf{1}, \mathbf{\bar{3}})$
$\hat{e}$	$\tilde{e}_R^*$	$e_R^*$	3	$(1, \mathbf{1}, \mathbf{1})$

- Superpotential:  $\mu$ -the only dimensional parameter.  $\mu$ -problem!

$$W_{\text{MSSM}} = \mu \hat{H}_u \hat{H}_d - Y_d \hat{d} \hat{q} \hat{H}_d - Y_e \hat{e} \hat{l} \hat{H}_d + Y_u \hat{u} \hat{q} \hat{H}_u$$



# MSSM: DM-nucleon scattering

Neutralino mass matrix:

$$m_{\tilde{\chi}^0} = \begin{pmatrix} M_1 & 0 & -\frac{1}{2}g_1 v_d & \frac{1}{2}g_1 v_u \\ 0 & M_2 & \frac{1}{2}g_2 v_d & -\frac{1}{2}g_2 v_u \\ -\frac{1}{2}g_1 v_d & \frac{1}{2}g_2 v_d & 0 & -\mu \\ \frac{1}{2}g_1 v_u & -\frac{1}{2}g_2 v_u & -\mu & 0 \end{pmatrix}$$

Choose eigenvalues as inputs can simplify calculation:

$$N_{i,j} = \frac{1}{\sqrt{C_i}} \begin{pmatrix} (\mu^2 - m_{\chi_i}^2) (M_2 - m_{\chi_i}) - M_Z^2 c_W^2 (m_{\chi_i} + 2\mu s_\beta c_\beta) \\ -M_Z^2 s_W c_W (m_{\chi_i} + 2\mu s_\beta c_\beta) \\ (M_2 - m_{\chi_i}) (m_{\chi_i} c_\beta + \mu s_\beta) M_Z s_W \\ -(M_2 - m_{\chi_i}) (m_{\chi_i} s_\beta + \mu c_\beta) M_Z s_W \end{pmatrix}_j$$

$C_i$ : Normalization factor

$$C_i = M_Z^2 c_W^2 (m_{\chi_i} + 2\mu s_\beta c_\beta) [M_Z^2 (m_{\chi_i} + 2\mu s_\beta c_\beta) + 2(\mu^2 - m_{\chi_i}^2) (m_{\chi_i} - M_2)] \\ + (m_{\chi_i} - M_2)^2 \left\{ M_Z^2 s_W^2 [(m_{\chi_i}^2 + \mu^2) + 4\mu m_{\chi_i} s_\beta c_\beta] + (m_{\chi_i}^2 - \mu^2)^2 \right\}$$

# MSSM: DM-nucleon scattering

DM is Bino-dominated.  $\sigma_{\tilde{\chi}-N}$  is given by:

$$C_{\tilde{\chi}_1^0 \tilde{\chi}_1^0 h} \simeq e \tan \theta_W \frac{m_Z}{\mu \left(1 - m_{\tilde{\chi}_1^0}^2 / \mu^2\right)} \left( \sin 2\beta + \frac{m_{\tilde{\chi}_1^0}}{\mu} \right)$$

$$C_{\tilde{\chi}_1^0 \tilde{\chi}_1^0 Z} \simeq \frac{e \tan \theta_W \cos 2\beta}{2} \frac{m_Z^2}{\mu^2 - m_{\tilde{\chi}_1^0}^2}$$

$$\sigma_{\tilde{\chi}_1^0-N}^{\text{SI}} \simeq 5 \times 10^{-45} \text{ cm}^2 \left( \frac{C_{\tilde{\chi}_1^0 \tilde{\chi}_1^0 h}}{0.1} \right)^2 \left( \frac{m_h}{125 \text{ GeV}} \right)^2$$

$$\sigma_{\tilde{\chi}_1^0-N}^{\text{SD}} \simeq 10^{-39} \text{ cm}^2 \left( \frac{C_{\tilde{\chi}_1^0 \tilde{\chi}_1^0 Z}}{0.1} \right)^2$$

- The expressions are valid only by assuming that  $|\mu| \gg |m_{\tilde{\chi}_1^0}|$ ;
- With the increase of  $|\mu|$ ,  $\sigma^{\text{SI}}$  and  $\sigma^{\text{SD}}$  decrease monotonously;
- Blind spot:  $\sin 2\beta + m_{\tilde{\chi}_1^0}/\mu = 0$  ( $\tan \beta = 1$ ) for SI (SD) scattering;
- For  $\tilde{B} - \tilde{H}$  co-annihilation,  $|m_{\tilde{\chi}_1^0}/\mu| \simeq 1$ ,  $\sigma^{\text{SI}}/\sigma^{\text{SD}}$  are enhanced.

## Computing with mass insertion method:

$$\begin{aligned}
 a_{\mu, \text{WHL}}^{\text{SUSY}} &= \frac{\alpha_2}{8\pi} \frac{m_\mu^2 \mu M_2 \tan \beta}{m_{\tilde{\nu}_\mu}^4} \left\{ 2f_C \left( \frac{M_2^2}{m_{\tilde{\nu}_\mu}^2}, \frac{\mu^2}{m_{\tilde{\nu}_\mu}^2} \right) - \frac{m_{\tilde{\nu}_\mu}^4}{m_{\tilde{\mu}_L}^4} f_N \left( \frac{M_2^2}{m_{\tilde{\mu}_L}^2}, \frac{\mu^2}{m_{\tilde{\mu}_L}^2} \right) \right\} \\
 a_{\mu, \text{BHL}}^{\text{SUSY}} &= \frac{\alpha_Y}{8\pi} \frac{m_\mu^2 \mu M_1 \tan \beta}{m_{\tilde{\mu}_L}^4} f_N \left( \frac{M_1^2}{m_{\tilde{\mu}_L}^2}, \frac{\mu^2}{m_{\tilde{\mu}_L}^2} \right) \\
 a_{\mu, \text{BHR}}^{\text{SUSY}} &= -\frac{\alpha_Y}{4\pi} \frac{m_\mu^2 \mu M_1 \tan \beta}{m_{\tilde{\mu}_R}^4} f_N \left( \frac{M_1^2}{m_{\tilde{\mu}_R}^2}, \frac{\mu^2}{m_{\tilde{\mu}_R}^2} \right) \\
 a_{\mu, \text{BLR}}^{\text{SUSY}} &= \frac{\alpha_Y}{4\pi} \frac{m_\mu^2 \mu M_1 \tan \beta}{M_1^4} f_N \left( \frac{m_{\tilde{\mu}_L}^2}{M_1^2}, \frac{m_{\tilde{\mu}_R}^2}{M_1^2} \right)
 \end{aligned}$$

where the loop functions are given by:

$$\begin{aligned}
 f_C(x, y) &= \frac{5 - 3(x + y) + xy}{(x - 1)^2(y - 1)^2} - \frac{2 \ln x}{(x - y)(x - 1)^3} + \frac{2 \ln y}{(x - y)(y - 1)^3} \\
 f_N(x, y) &= \frac{-3 + x + y + xy}{(x - 1)^2(y - 1)^2} + \frac{2x \ln x}{(x - y)(x - 1)^3} - \frac{2y \ln y}{(x - y)(y - 1)^3}
 \end{aligned}$$

**Very good approximation!**

- The following processes are considered:

$$pp \rightarrow \tilde{\chi}_i^0 \tilde{\chi}_j^\pm, \quad i = 2, 3, 4; \quad j = 1, 2$$

$$pp \rightarrow \tilde{\chi}_i^\pm \tilde{\chi}_j^\mp, \quad i = 1, 2; \quad j = 1, 2$$

$$pp \rightarrow \tilde{\chi}_i^0 \tilde{\chi}_j^0, \quad i = 2, 3, 4; \quad j = 2, 3, 4$$

$$pp \rightarrow \tilde{\mu}_i \tilde{\mu}_j, \tilde{\nu} \tilde{\nu} \quad i = L, R; \quad j = L, R$$

- All LHC searches for electroweakinos and sleptons are considered, a total of 14 analyses for Run-II data.
- Newly added important analyses:
  - ① ATLAS search for 3 lepton plus missing  $E_T$  signal, see CERN-EP-2021-059, or arXiv: 2106.01676.
  - ② CMS search for 2 lepton plus missing  $E_T$  signal, arXiv: 2012.08600.
- Other important constraints:  
CERN-EP-2019-106/2019-263/2019-188, CMS-SUS-17-004/20-001.

# MSSM: LHC search for SUSY

Analysis	Simplified Scenario	Signal of Final State	Luminosity
<b>CMS-SUS-17-010</b> (arXiv:1807.07799)	$\tilde{\chi}_1^\pm \tilde{\chi}_1^\mp \rightarrow W^\pm \tilde{\chi}_1^0 W^\mp \tilde{\chi}_1^0$ $\tilde{\chi}_1^\pm \tilde{\chi}_1^\mp \rightarrow \nu \bar{\ell} / \ell \bar{\nu} \rightarrow \ell \ell \nu \tilde{\chi}_1^0 \tilde{\chi}_1^0$	$2\ell + E_T^{\text{miss}}$	35.9 fb <sup>-1</sup>
<b>CMS-SUS-17-009</b> (arXiv:1806.05264)	$\tilde{\ell} \bar{\ell} \rightarrow \ell \ell \tilde{\chi}_1^0 \tilde{\chi}_1^0$	$2\ell + E_T^{\text{miss}}$	35.9 fb <sup>-1</sup>
<b>CMS-SUS-17-004</b> (arXiv:1801.03957)	$\tilde{\chi}_2^0 \tilde{\chi}_1^\pm \rightarrow Wh(Z) \tilde{\chi}_1^0 \tilde{\chi}_1^0$	$n\ell(\geq 0) + nj(\geq 0) + E_T^{\text{miss}}$	35.9 fb <sup>-1</sup>
<b>CMS-SUS-16-045</b> (arXiv:1709.00384)	$\tilde{\chi}_2^0 \tilde{\chi}_1^\pm \rightarrow W^\pm \tilde{\chi}_1^0 h \tilde{\chi}_1^0$	$1\ell 2b + E_T^{\text{miss}}$	35.9 fb <sup>-1</sup>
<b>CMS-SUS-16-039</b> (arxiv:1709.05406)	$\tilde{\chi}_2^0 \tilde{\chi}_1^\pm \rightarrow \ell \nu \bar{\ell} \bar{\ell}$ $\tilde{\chi}_2^0 \tilde{\chi}_1^\pm \rightarrow \bar{\tau} \nu \bar{\ell} \bar{\ell}$ $\tilde{\chi}_2^0 \tilde{\chi}_1^\pm \rightarrow \bar{\tau} \nu \bar{\tau} \tau$ $\tilde{\chi}_2^0 \tilde{\chi}_1^\pm \rightarrow WZ \tilde{\chi}_1^0 \tilde{\chi}_1^0$ $\tilde{\chi}_2^0 \tilde{\chi}_1^\pm \rightarrow WH \tilde{\chi}_1^0 \tilde{\chi}_1^0$	$n\ell(\geq 0)(\tau) + E_T^{\text{miss}}$	35.9 fb <sup>-1</sup>
<b>CMS-SUS-16-034</b> (arXiv:1709.08908)	$\tilde{\chi}_2^0 \tilde{\chi}_1^\pm \rightarrow W \tilde{\chi}_1^0 Z(h) \tilde{\chi}_1^0$	$n\ell(\geq 2) + nj(\geq 1) E_T^{\text{miss}}$	35.9 fb <sup>-1</sup>
<b>CERN-EP-2017-303</b> (arXiv:1803.02762)	$\tilde{\chi}_2^0 \tilde{\chi}_1^\pm \rightarrow WZ \tilde{\chi}_1^0 \tilde{\chi}_1^0$ $\tilde{\chi}_1^\pm \tilde{\chi}_1^\mp \rightarrow \nu \bar{\ell} / \ell \bar{\nu} \rightarrow \ell \ell \nu \tilde{\chi}_1^0 \tilde{\chi}_1^0$ $\bar{\ell} \bar{\ell} \rightarrow \ell \ell \tilde{\chi}_1^0 \tilde{\chi}_1^0$	$n\ell(\geq 2) + E_T^{\text{miss}}$	35.9 fb <sup>-1</sup>
<b>CERN-EP-2018-306</b> (arXiv:1812.09432)	$\tilde{\chi}_2^0 \tilde{\chi}_1^\pm \rightarrow Wh \tilde{\chi}_1^0 \tilde{\chi}_1^0$	$n\ell(\geq 0) + nj(\geq 0) + nb(\geq 0) + n\gamma(\geq 0) + E_T^{\text{miss}}$	35.9 fb <sup>-1</sup>
<b>CERN-EP-2018-113</b> (arXiv:1806.02293)	$\tilde{\chi}_2^0 \tilde{\chi}_1^\pm \rightarrow WZ \tilde{\chi}_1^0 \tilde{\chi}_1^0$	$n\ell(\geq 2) + nj(\geq 0) + E_T^{\text{miss}}$	35.9 fb <sup>-1</sup>
<b>CERN-EP-2019-263</b> (arXiv:1912.08479)	$\tilde{\chi}_2^0 \nu \tilde{\chi}_1^\pm \rightarrow W \tilde{\chi}_1^0 Z \tilde{\chi}_1^0 \rightarrow \ell \nu \ell \tilde{\chi}_1^0 \tilde{\chi}_1^0$	$3\ell + E_T^{\text{miss}}$	139 fb <sup>-1</sup>
<b>CERN-EP-2019-106</b> (arXiv:1908.08215)	$\bar{\ell} \bar{\ell} \rightarrow \ell \ell \tilde{\chi}_1^0 \tilde{\chi}_1^0$ $\tilde{\chi}_1^\pm \tilde{\chi}_1^\mp \rightarrow \nu \bar{\ell} / \ell \bar{\nu} \rightarrow \ell \ell \nu \tilde{\chi}_1^0 \tilde{\chi}_1^0$	$2\ell + E_T^{\text{miss}}$	139 fb <sup>-1</sup>
<b>CERN-EP-2019-188</b> (arXiv:1909.09226)	$\tilde{\chi}_2^0 \tilde{\chi}_1^\pm \rightarrow Wh \tilde{\chi}_1^0 \tilde{\chi}_1^0$	$1\ell + h(\rightarrow b\bar{b}) + E_T^{\text{miss}}$	139 fb <sup>-1</sup>

Table 1: Signal of final state for electroweakino pair-production processes.

# MSSM: Numerical results

## Preferred DM annihilation channels:

### Co-annihilating with a Wino-dominated NLSP:

- ① Constraints on LHC search for SUSY is relatively weak.
- ② Tri-lepton signal for compressed spectrum:  $|m_{\tilde{\chi}_1^0}| > 210$  GeV.

### Co-annihilating with Sletpon NLSP:



- ① The Slepton may be right-handed or left-handed.
- ② LHC cosntraints are very strong!

## Excluded annihilation channels:

### Co-annihilating with a Higgsino-dominated NLSP:

- ① DM direct detection experiments prefer an excessively large  $|\mu|$ .
- ② Unable to explain the muon g-2 anomaly since  $|\mu|$  is large.

### Resonant $h/Z$ annihilation: tight LHC constraints.



- Based on more than  $10^8$  samplings and more than  $10^6$  simulations;
- $|m_{\tilde{\chi}_1^0}| \gtrsim 210$  GeV:  $3\ell + E_T^{\text{Miss}}$  signal for compressed spectrum;
- DM direct detection experiments:  $|\mu| \gtrsim 300$  GeV;
- DM +  $a_\mu$  + LHC:  $|\mu| \gtrsim 450$  GeV,  $\sim 1\%$  tuning for  $m_Z$ .

## Section IV

### Example II: $Z_3$ -MSSM



# $Z_3$ -NMSSM: Theoretical preliminaries

- Field content and gauge group

SF	Spin 0	Spin $\frac{1}{2}$	Generations	$(U(1) \otimes SU(2) \otimes SU(3))$
$\hat{q}$	$\tilde{q}$	$q$	3	$(\frac{1}{6}, \mathbf{2}, \mathbf{3})$
$\hat{l}$	$\tilde{l}$	$l$	3	$(-\frac{1}{2}, \mathbf{2}, \mathbf{1})$
$\hat{H}_d$	$H_d$	$\tilde{H}_d$	1	$(-\frac{1}{2}, \mathbf{2}, \mathbf{1})$
$\hat{H}_u$	$H_u$	$\tilde{H}_u$	1	$(\frac{1}{2}, \mathbf{2}, \mathbf{1})$
$\hat{d}$	$\tilde{d}_R^*$	$d_R^*$	3	$(\frac{1}{3}, \mathbf{1}, \overline{\mathbf{3}})$
$\hat{u}$	$\tilde{u}_R^*$	$u_R^*$	3	$(-\frac{2}{3}, \mathbf{1}, \overline{\mathbf{3}})$
$\hat{e}$	$\tilde{e}_R^*$	$e_R^*$	3	$(1, \mathbf{1}, \mathbf{1})$
$\hat{S}$	$S$	$\tilde{S}$	1	$(0, \mathbf{1}, \mathbf{1})$

- Superpotential: Try to solve  $\mu$ - and little hierarchy problems of the MSSM. There is no dimensional parameters in the superpotential. However, once  $Z_3$ -symmetry was spontaneously broken, domain wall problem and tadpole problem will be induced! Occam's razor was used incorrectly.

$$W_{\text{NMSSM}} = W_{\text{Yukawa}} + \lambda \hat{S} \hat{H}_u \hat{H}_d + \frac{1}{3} \kappa \hat{S}^3$$

- Neutralino mass matrix

$$\mathcal{M} = \begin{pmatrix} M_1 & 0 & -\frac{g_1 v_d}{\sqrt{2}} & \frac{g_1 v_u}{\sqrt{2}} & 0 \\ & M_2 & \frac{g_2 v_d}{\sqrt{2}} & -\frac{g_2 v_u}{\sqrt{2}} & 0 \\ & & 0 & -\mu_{eff} & -\lambda v_u \\ & & & 0 & -\lambda v_d \\ & & & & \frac{2\kappa}{\lambda} \mu_{eff} \end{pmatrix}$$

- DM may be Singlino-dominated. Its mass and couplings are given by

$$m_{\tilde{\chi}_1^0} \simeq \frac{2\kappa}{\lambda} \mu + \frac{\lambda^2 v^2}{\mu^2} (\mu \sin 2\beta - \frac{2\kappa}{\lambda} \mu), \quad C_{\tilde{\chi}_1^0 \tilde{\chi}_1^0 Z} \simeq \frac{m_Z}{\sqrt{2}v} \left( \frac{\lambda v}{\mu_{eff}} \right)^2 \frac{\cos 2\beta}{1 - (m_{\tilde{\chi}_1^0}/\mu_{eff})^2},$$

$$C_{\tilde{\chi}_1^0 \tilde{\chi}_1^0 h_i} \simeq \sqrt{2} \lambda \left( \frac{\lambda v}{\mu_{eff}} \right) \frac{V_{hi}^{SM} (m_{\tilde{\chi}_1^0}/\mu_{eff} - \sin 2\beta)}{1 - (m_{\tilde{\chi}_1^0}/\mu_{eff})^2} - \sqrt{2} \lambda \left( \frac{\lambda v}{\mu_{eff}} \right) \frac{V_{hi}^{NSM} \cos 2\beta}{1 - (m_{\tilde{\chi}_1^0}/\mu_{eff})^2}$$

$$+ \lambda \left( \frac{\lambda v}{\mu_{eff}} \right)^2 \frac{V_{hi}^S \sin 2\beta}{\sqrt{2} [1 - (m_{\tilde{\chi}_1^0}/\mu_{eff})^2]} - \sqrt{2} \kappa V_{hi}^S \left[ 1 + \left( \frac{\lambda v}{\mu_{eff}} \right)^2 \frac{2}{1 - (m_{\tilde{\chi}_1^0}/\mu_{eff})^2} \right],$$

# $Z_3$ -NMSSM: DM-nucleon scattering

$$\sigma_{\tilde{\chi}_1^0-N}^{\text{SI}} \simeq 5 \times 10^{-45} \text{cm}^2 \times \left( \frac{\mathcal{A}}{0.1} \right)^2,$$

$$\sigma_{\tilde{\chi}_1^0-N}^{\text{SD}} \simeq 10^{-39} \text{cm}^2 \left( \frac{C_{\tilde{\chi}_1^0 \tilde{\chi}_1^0 Z}}{0.1} \right)^2,$$

$$\mathcal{A} \simeq \left( \frac{125 \text{GeV}}{m_h} \right)^2 V_h^{\text{SM}} C_{\tilde{\chi}_1^0 \tilde{\chi}_1^0 h} + \left( \frac{125 \text{GeV}}{m_{h_s}} \right)^2 V_{h_s}^{\text{SM}} C_{\tilde{\chi}_1^0 \tilde{\chi}_1^0 h_s}$$

$$\simeq \left( \frac{125 \text{GeV}}{m_h} \right)^2 \lambda \frac{\sqrt{2} \lambda v}{\mu_{eff}} \frac{(m_{\tilde{\chi}_1^0} / \mu_{eff} - \sin 2\beta)}{1 - (\mathbf{m}_{\tilde{\chi}_1^0} / \mu_{eff})^2},$$

$$C_{\tilde{\chi}_1^0 \tilde{\chi}_1^0 Z} \simeq \frac{m_Z}{\sqrt{2}v} \left( \frac{\lambda v}{\mu_{eff}} \right)^2 \frac{\cos 2\beta}{1 - (\mathbf{m}_{\tilde{\chi}_1^0} / \mu_{eff})^2}.$$

- DM properties are described by four independent parameters:

$$\lambda, \quad \mu_{eff}, \quad m_{\tilde{\chi}_1^0}, \quad \tan \beta.$$

- $m_{\tilde{\chi}_1^0}$  and  $\kappa$  are correlated,  $\lambda$  and  $\kappa$  are correlated by  $2\kappa / \lambda < 1$ ;
- $\lambda \lesssim 0.1$  is preferred to suppress DM-nucleon scattering.

# $Z_3$ -NMSSM: Dominant annihilation channels

Conditions to obtain the measured DM abundance:

- ❶  $\tilde{\chi}_1^0 \tilde{\chi}_1^0 \rightarrow t\bar{t}$ :  $s$ -channel exchange of  $Z$  and Higgs bosons.

$$|C_{\tilde{\chi}_1^0 \tilde{\chi}_1^0 G^0}| = \frac{\sqrt{2} m_{\tilde{\chi}_1^0}}{v} \left( \frac{\lambda v}{\mu_{eff}} \right)^2 \frac{\cos 2\beta}{1 - (m_{\tilde{\chi}_1^0}/\mu_{eff})^2} \simeq 0.1.$$

- ❷  $\tilde{\chi}_1^0 \tilde{\chi}_1^0 \rightarrow h_s A_s$ :  $s$ -channel exchange of Higgs bosons,  $t$ -channel exchange of neutralinos.

$$|C_{\tilde{\chi}_1^0 \tilde{\chi}_1^0 h_s}| = |C_{\tilde{\chi}_1^0 \tilde{\chi}_1^0 A_s}| = -\sqrt{2} \kappa \simeq 0.2 \times \left( \frac{m_{\tilde{\chi}_1^0}}{300 \text{ GeV}} \right)^{1/2}.$$

- ❸  $\tilde{\chi}_1^0 \tilde{\chi}_1^0 \rightarrow h A_s$ :  $s$ -channel exchange of Higgs bosons,  $t$ -channel exchange of neutralinos.

$$\lambda^3 \sin 2\beta \simeq \left( \frac{\mu_{eff}}{700 \text{ GeV}} \right)^2.$$

$\lambda > 0.3$  is preferred to predict the measured abundance.

# $Z_3$ -NMSSM: Summary

## The status of $Z_3$ -NMSSM:

Model	MSSM	$Z_3$ -NMSSM		GNMSSM	Type-I NMSSM	B-L NMSSM
DM candidate	Bino	Bino	Singlino	Singlino	Sneutrino	Bino, Singlino, BLino, Bileptino, and sneutrino
DM physics	×	×	×	✓	✓	✓
LHC and $\Delta a_\mu$	×	×	×	✓	✓	✓
EWSB	×	×	✓	✓	✓	✓
Neutrino	×	×	×	×	✓	✓
Higgs mass	×	×	×	×	✓	✓

The singlino-dominated DM scenario in  $Z_3$ -NMSSM has been tightly limited. The phenomenology of the bino-dominated DM scenario is roughly same as that of the MSSM.

## Part IV

### Example III: General MSSM

# GNMSSM: Motivation and superpotential

- Chiral Superfields

SF	Spin 0	Spin $\frac{1}{2}$	Generations	$(U(1) \otimes SU(2) \otimes SU(3))$
$\hat{q}$	$\tilde{q}$	$q$	3	$(\frac{1}{6}, \mathbf{2}, \mathbf{3})$
$\hat{l}$	$\tilde{l}$	$l$	3	$(-\frac{1}{2}, \mathbf{2}, \mathbf{1})$
$\hat{H}_d$	$H_d$	$\tilde{H}_d$	1	$(-\frac{1}{2}, \mathbf{2}, \mathbf{1})$
$\hat{H}_u$	$H_u$	$\tilde{H}_u$	1	$(\frac{1}{2}, \mathbf{2}, \mathbf{1})$
$\hat{d}$	$\tilde{d}_R^*$	$d_R^*$	3	$(\frac{1}{3}, \mathbf{1}, \mathbf{\bar{3}})$
$\hat{u}$	$\tilde{u}_R^*$	$u_R^*$	3	$(-\frac{2}{3}, \mathbf{1}, \mathbf{\bar{3}})$
$\hat{e}$	$\tilde{e}_R^*$	$e_R^*$	3	$(1, \mathbf{1}, \mathbf{1})$
$\hat{s}$	$S$	$\bar{S}$	1	$(0, \mathbf{1}, \mathbf{1})$

- Superpotential

$$W_{\text{GNMSSM}} = W_{\text{Yukawa}} + \lambda \hat{S} \hat{H}_u \cdot \hat{H}_d + \frac{1}{3} \kappa \hat{S}^3 + \mu \hat{H}_u \cdot \hat{H}_d + \frac{1}{2} \mu' \hat{S}^2 + \xi \hat{S}$$

- 1 Solve domain wall and tadpole problems in  $Z_3$ -NMSSM.
- 2  $Z_3$ -violating terms from an underlying theory with  $Z_4^n$  or  $Z_8^n$  symmetry.

## CP-even Higgs mass matrix

$$\begin{aligned}\mathcal{M}_{S,11}^2 &= \frac{2 [\mu_{eff}(\lambda A_\lambda + \kappa \mu_{eff} + \lambda \mu') + \lambda m_3^2]}{\lambda \sin 2\beta} + \frac{1}{2}(2m_Z^2 - \lambda^2 v^2) \sin^2 2\beta, \\ \mathcal{M}_{S,12}^2 &= -\frac{1}{4}(2m_Z^2 - \lambda^2 v^2) \sin 4\beta, \\ \mathcal{M}_{S,13}^2 &= -\frac{1}{\sqrt{2}}(\lambda A_\lambda + 2\kappa \mu_{eff} + \lambda \mu') v \cos 2\beta, \\ \mathcal{M}_{S,22}^2 &= m_Z^2 \cos^2 2\beta + \frac{1}{2}\lambda^2 v^2 \sin^2 2\beta, \\ \mathcal{M}_{S,23}^2 &= \frac{v}{\sqrt{2}} [2\lambda(\mu_{eff} + \mu) - (\lambda A_\lambda + 2\kappa \mu_{eff} + \lambda \mu') \sin 2\beta], \\ \mathcal{M}_{S,33}^2 &= \frac{\lambda(A_\lambda + \mu') \sin 2\beta}{4\mu_{eff}} \lambda v^2 + \frac{\mu_{eff}}{\lambda} (\kappa A_\kappa + \frac{4\kappa^2 \mu_{eff}}{\lambda} + 3\kappa \mu') - \frac{\mu}{2\mu_{eff}} \lambda^2 v^2, \\ m_{h_s}^2 &\simeq \mathcal{M}_{S,33}^2 - \frac{\mathcal{M}_{S,13}^4}{\mathcal{M}_{S,11}^2 - \mathcal{M}_{S,33}^2} \\ &\simeq \frac{\mu_{eff}}{\lambda} (\kappa A_\kappa + \frac{4\kappa^2 \mu_{eff}}{\lambda} + 3\kappa \mu') - \frac{\mu_{tot}}{2\mu_{eff}} \lambda^2 v^2 + \frac{1}{2} \lambda^2 v^2,\end{aligned}$$

$m_{h_s}$  can be modified freely by  $\mu'$ .



## CP-odd Higgs mass matrix

$$\begin{aligned}\mathcal{M}_{P,11}^2 &= \frac{2[\mu_{eff}(\lambda A_\lambda + \kappa\mu_{eff} + \lambda\mu') + \lambda m_3^2]}{\lambda \sin 2\beta}, \\ \mathcal{M}_{P,22}^2 &= \frac{(\lambda A_\lambda + 4\kappa\mu_{eff} + \lambda\mu') \sin 2\beta}{4\mu_{eff}} \lambda v^2 - \frac{\kappa\mu_{eff}}{\lambda} (3A_\kappa + \mu') - \frac{\mu}{2\mu_{eff}} \lambda^2 v^2 - 2m_S'^2, \\ \mathcal{M}_{P,12}^2 &= \frac{v}{\sqrt{2}} (\lambda A_\lambda - 2\kappa\mu_{eff} - \lambda\mu').\end{aligned}\tag{1}$$

$$\begin{aligned}m_{A_s}^2 &\simeq \mathcal{M}_{P,22}^2 - \frac{\mathcal{M}_{P,12}^4}{\mathcal{M}_{P,11}^2 - \mathcal{M}_{P,22}^2} \\ &\simeq -\frac{\kappa\mu_{eff}}{\lambda} (3A_\kappa + \mu') - \frac{\mu_{tot}}{2\mu_{eff}} \lambda^2 v^2 + \frac{1}{2} \lambda^2 v^2 - 2m_S'^2.\end{aligned}\tag{2}$$

$m_{A_s}$  can be modified freely by  $m_S'$ .

# GNMSSM: DM properties

- Neutralino mass matrix

$$m_{\tilde{\chi}_i^0} = \begin{pmatrix} M_1 & 0 & -\frac{1}{2}g_1 v_d & \frac{1}{2}g_1 v_u & 0 \\ 0 & M_2 & \frac{1}{2}g_2 v_d & -\frac{1}{2}g_2 v_u & 0 \\ -\frac{1}{2}g_1 v_d & \frac{1}{2}g_2 v_d & 0 & -\mu - \mu_{\text{eff}} & -\frac{1}{\sqrt{2}}v_u \lambda \\ \frac{1}{2}g_1 v_u & -\frac{1}{2}g_2 v_u & -\mu - \mu_{\text{eff}} & 0 & -\frac{1}{\sqrt{2}}v_d \lambda \\ 0 & 0 & -\frac{1}{\sqrt{2}}v_u \lambda & -\frac{1}{\sqrt{2}}v_d \lambda & \frac{2\kappa}{\lambda}\mu_{\text{eff}} + \mu' \end{pmatrix}$$

Mass and couplings of the singlino-dominated DM are given by:

$$m_{\tilde{\chi}_1^0} \simeq \frac{2\kappa}{\lambda}\mu_{\text{eff}} + \frac{1}{2} \frac{\lambda^2 v^2 (m_{\tilde{\chi}_1^0} - \mu_{\text{tot}} \sin 2\beta)}{m_{\tilde{\chi}_1^0}^2 - \mu_{\text{tot}}^2},$$

$$C_{\tilde{\chi}_1^0 \tilde{\chi}_1^0 h_i} = C_{\tilde{\chi}_1^0 \tilde{\chi}_1^0 h_i}^{\text{Z}_3\text{-NMSSM}}|_{\mu_{\text{eff}} \rightarrow \mu_{\text{tot}}},$$

$$C_{\tilde{\chi}_1^0 \tilde{\chi}_1^0 Z} = C_{\tilde{\chi}_1^0 \tilde{\chi}_1^0 Z}^{\text{Z}_3\text{-NMSSM}}|_{\mu_{\text{eff}} \rightarrow \mu_{\text{tot}}}.$$

- DM mass and  $\kappa$  are not correlated,  $\lambda$  and  $\kappa$  are not correlated!**
- DM properties are described by **five** independent parameters:  
 $m_{\tilde{\chi}_1^0}$ ,  $\lambda$ ,  $\kappa$ ,  $\tan\beta$ , and  $\mu_{\text{tot}} \equiv \mu + \mu_{\text{eff}}$ .
- $\lambda \lesssim 0.1$  is preferred to suppress DM-nucleon scatterings.

# GNMSSM: Dominant annihilation channels

## Conditions to obtain the measured DM abundance:

- ❶  $\tilde{\chi}_1^0 \tilde{\chi}_1^0 \rightarrow t\bar{t}$ :  $s$ -channel exchange of  $Z$  and Higgs bosons.

$$|C_{\tilde{\chi}_1^0 \tilde{\chi}_1^0 G^0}| = \frac{\sqrt{2} m_{\tilde{\chi}_1^0}}{v} \left( \frac{\lambda v}{\mu_{tot}} \right)^2 \frac{\cos 2\beta}{1 - (m_{\tilde{\chi}_1^0}/\mu_{tot})^2} \simeq 0.1.$$

- ❷  $\tilde{\chi}_1^0 \tilde{\chi}_1^0 \rightarrow h_s A_s$ :  $s$ -channel exchange of Higgs bosons,  $t$ -channel exchange of neutralinos.

$$|C_{\tilde{\chi}_1^0 \tilde{\chi}_1^0 h_s}| = |C_{\tilde{\chi}_1^0 \tilde{\chi}_1^0 A_s}| = -\sqrt{2} \kappa \simeq 0.2 \times \left( \frac{m_{\tilde{\chi}_1^0}}{300 \text{ GeV}} \right)^{1/2}.$$

- ❸  $\tilde{\chi}_1^0 \tilde{\chi}_1^0 \rightarrow h A_s$ :  $s$ -channel exchange of Higgs bosons,  $t$ -channel exchange of neutralinos.

$$\lambda^3 \sin 2\beta \simeq \left( \frac{\mu}{700 \text{ GeV}} \right)^2.$$

**Singlet-dominated particles may form a secluded DM sector:**  
measured abundance obtained by  $\tilde{\chi}_1^0 \tilde{\chi}_1^0 \rightarrow h_s A_s$  (via adjusting  $\kappa$ );  
DM-nucleon scatterings suppressed by a small  $\lambda v/\mu_{tot}$ .  
**The simplest SUSY framework to realize secluded DM sector.**

# GNMSSM: Dominant annihilation channels

$h \equiv h_1$ scenario: $\ln Z = -65.79 \pm 0.046$			
$\tilde{\chi}_1^0 \tilde{\chi}_1^0 \rightarrow h_s A_s$	$\tilde{\chi}_1^0 \tilde{\chi}_1^0 \rightarrow t \bar{t}$	$\tilde{\chi}_1^0 \tilde{\chi}_1^0 \rightarrow h_s h_s$	Co-annihilation
88%	8%	3%	0.7%
$h \equiv h_2$ scenario: $\ln Z = -68.23 \pm 0.051$			
$\tilde{\chi}_1^0 \tilde{\chi}_1^0 \rightarrow h_s A_s$	$\tilde{\chi}_1^0 \tilde{\chi}_1^0 \rightarrow t \bar{t}$	Co-annihilation	$h$ -funnel
76%	12%	11.6%	0.3%

**Table 2:** Dominant annihilation channels and their normalized posterior probabilities for  $h \equiv h_1$  and  $h \equiv h_2$  scenarios. In obtaining the values in this table, each sample's most critical channel for the abundance was identified and sequentially used to classify the samples. The posterior probability densities of the same type of samples were then summed.

**$\tilde{\chi}_1^0 \tilde{\chi}_1^0 \rightarrow h_s A_s$  always played a role in DM annihilation.**

# GNMSSM: Explaining the Muon $g-2$ anomaly

## Characteristics:

- ① Roughly same loop contributions as the MSSM.
- ② **DM physics is changed.**
- ③ **LHC constraints is alleviated significantly.**
- ④ **Vacuum becomes more stable.**

## Mechanism to alleviate the LHC constraints:

- ① DM must be heavy to achieve the measured relic density.
- ② For singlino-dominated DM, heavy sparticles prefer to decay into NLSP or NNLSP first; their decay chains are lengthened.
- ③ Light singlet Higgs bosons may act as the sparticle decay products.

# GNMSSM: Current Status

Model	MSSM	Z <sub>3</sub> -NMSSM		GNMSSM	Type-I/ISS-NMSSM	B-L NMSSM
DM candidate	Bino	Bino	Singlino	Singlino	Sneutrino	Bino, Singlino, BLino, Bileptino, and sneutrino
DM physics	×	×	×	✓	✓	✓
LHC and $\Delta a_\mu$	×	×	×	✓	✓	✓
EWSB	×	×	✓	✓	✓	✓
Neutrino	×	×	×	×	✓	✓
Higgs mass	×	×	×	×	✓	✓

## Part Five

### Example IV: Type-I MSSM

# Type-I NMSSM

- Chiral Superfields

SF	Spin 0	Spin $\frac{1}{2}$	Generations	$(U(1) \otimes SU(2) \otimes SU(3))$
$\hat{q}$	$\tilde{q}$	$q$	3	$(\frac{1}{6}, \mathbf{2}, \mathbf{3})$
$\hat{l}$	$\tilde{l}$	$l$	3	$(-\frac{1}{2}, \mathbf{2}, \mathbf{1})$
$\hat{H}_d$	$H_d$	$\tilde{H}_d$	1	$(-\frac{1}{2}, \mathbf{2}, \mathbf{1})$
$\hat{H}_u$	$H_u$	$\tilde{H}_u$	1	$(\frac{1}{2}, \mathbf{2}, \mathbf{1})$
$\hat{d}$	$\tilde{d}_R^*$	$d_R^*$	3	$(\frac{1}{3}, \mathbf{1}, \bar{\mathbf{3}})$
$\hat{u}$	$\tilde{u}_R^*$	$u_R^*$	3	$(-\frac{2}{3}, \mathbf{1}, \bar{\mathbf{3}})$
$\hat{e}$	$\tilde{e}_R^*$	$e_R^*$	3	$(1, \mathbf{1}, \mathbf{1})$
$\hat{\nu}$	$\tilde{\nu}_R^*$	$\nu_R^*$	3	$(0, \mathbf{1}, \mathbf{1})$
$\hat{s}$	$S$	$\tilde{S}$	1	$(0, \mathbf{1}, \mathbf{1})$

- Superpotential

$$W_{\text{Type-I}} = W_{\text{Yukawa}} + \lambda \hat{S} \hat{H}_u \cdot \hat{H}_d + \frac{1}{3} \kappa \hat{S}^3 + \bar{\lambda}_\nu \hat{s} \hat{\nu} \hat{\nu} + Y_\nu \hat{l} \cdot \hat{H}_u \hat{\nu}$$

- 1 Provide mechanisms to generate neutrino mass and mixing, and leptogenesis.
- 2 Lightest sneutrino may act as a feasible DM candidate.



- Sneutrino mass matrix

$$\mathcal{M}_{\tilde{\nu}}^2 = \begin{pmatrix} m_{L\bar{L}}^2 & \frac{m_{LR}^2 + m_{L\bar{R}}^2 + \text{c.c.}}{2} & 0 & i \frac{m_{LR}^2 - m_{L\bar{R}}^2 - \text{c.c.}}{2} \\ \frac{m_{LR}^2 + m_{L\bar{R}}^2 + \text{c.c.}}{2} & m_{R\bar{R}}^2 + m_{RR}^2 + m_{RR}^{2*} & i \frac{m_{LR}^2 - m_{L\bar{R}}^2 - \text{c.c.}}{2} & i (m_{RR}^2 - m_{RR}^{2*}) \\ 0 & i \frac{m_{LR}^2 - m_{L\bar{R}}^2 - \text{c.c.}}{2} & m_{L\bar{L}}^2 & \frac{-m_{LR}^2 + m_{L\bar{R}}^2 + \text{c.c.}}{2} \\ i \frac{m_{LR}^2 - m_{L\bar{R}}^2 - \text{c.c.}}{2} & i (m_{RR}^2 - m_{RR}^{2*}) & \frac{-m_{LR}^2 + m_{L\bar{R}}^2 + \text{c.c.}}{2} & m_{R\bar{R}}^2 - m_{RR}^2 - m_{RR}^{2*} \end{pmatrix}$$

- Chiral mixing can be neglected. DM may be purely right-handed sneutrino.
- Lepton number violating interactions split right-handed CP-even and CP-odd sneutrinos.

- Expression of DM-nucleon scattering cross section:

$$\sigma_{\tilde{\nu}_1-N}^{\text{SI}} \simeq 4.2 \times 10^{-44} \text{ cm}^2 \times \left( \frac{125 \text{ GeV}}{m_h} \right)^4 \times \left( \frac{C_{\tilde{\nu}_1^* \tilde{\nu}_1} \text{Re}[S]}{m_{\tilde{\nu}_1}} \times \delta \sin \theta \cos \theta - \frac{\cos \beta C_{\tilde{\nu}_1^* \tilde{\nu}_1} \text{Re}[H_d^0] + \sin \beta C_{\tilde{\nu}_1^* \tilde{\nu}_1} \text{Re}[H_u^0]}{m_{\tilde{\nu}_1}} \times (1 + \delta \sin^2 \theta) \right)^2$$

where

$$C_{\tilde{\nu}_1 \tilde{\nu}_1 h_i} = \frac{\lambda \lambda_\nu M_W}{g} (\sin \beta Z_{i1} + \cos \beta Z_{i2}) - \left[ \frac{\sqrt{2}}{\lambda} (2\lambda_\nu^2 + \kappa \lambda_\nu) \mu - \frac{\lambda_\nu A_{\lambda_\nu}}{\sqrt{2}} \right] Z_{i3},$$

$$\delta = m_h^2 / m_{h_s}^2 - 1.$$

**DM-nucleon scatterings are naturally suppressed!**  
**A general conclusion for singlet-dominated DM.**

## DM annihilation channels

- $\tilde{\nu}_1 \tilde{H} \rightarrow XY, \tilde{H} \tilde{H}' \rightarrow X'Y'$ :  
 $m_{\tilde{\nu}_1} \simeq \mu$ , co-annihilate with Higgsino-dominated electroweakinos.
- $\tilde{\nu}_1 \tilde{\nu}_1 \rightarrow SS^*$ :  
s-channel Higgs exchange, t/u-channel sneutrino exchange, and a four-point interaction.
- $\tilde{\nu}_1 \tilde{\nu}_1 \rightarrow \nu_R \bar{\nu}_R$ :  
s-channel Higgs exchange and t/u-channel neutralino exchange.
- $\tilde{\nu}_1 \tilde{\nu}_1 \rightarrow VV^*, VS, f\bar{f}$ :  
s-channel Higgs exchange.

In the DM annihilation processes, the singlet field as a propagator and final states contributing the most to the correct residual density of dark matter.

CP-even light $h_s$ scenario: $\ln Z = -40.7 \pm 0.20$			
Annihilation characteristics		Percent	
Coannihilation	$\tilde{\nu}_1 \chi_1 \rightarrow XY$	37%	38.2%
	$\tilde{\nu}_1 \tilde{\nu}_1^I \rightarrow \nu_4 \nu_4$	1.1%	
	$\tilde{\nu}_1^I \tilde{\nu}_1^I \rightarrow h_s h_s$	0.1%	
Secluded DM sector	$\tilde{\nu}_1 \tilde{\nu}_1 \rightarrow A_s A_s$	0.2%	55.3%
	$\tilde{\nu}_1 \tilde{\nu}_1 \rightarrow h_s h_s$	54%	
	$\tilde{\nu}_1 \tilde{\nu}_1 \rightarrow \nu_4 \nu_4$	1.1%	
Higgs portal	$\tilde{\nu}_1 \tilde{\nu}_1 \rightarrow h h_s$	0.2%	6.5%
	$\tilde{\nu}_1 \tilde{\nu}_1 \rightarrow h h$	0.3%	
	$\tilde{\nu}_1 \tilde{\nu}_1 \rightarrow g g$	0.1%	
	$\tilde{\nu}_1 \tilde{\nu}_1 \rightarrow b \bar{b}$	1.8%	
	$\tilde{\nu}_1 \tilde{\nu}_1 \rightarrow W^+ W^-$	4.1%	

**Table 3:** The annihilation mechanisms and channels in CP-even light  $h_s$  scenario, where  $\chi_1 \equiv \{\tilde{\chi}_1^\pm, \tilde{\chi}_1^0, \tilde{\chi}_2^0\}$ , X and Y represent any possible final states, and  $\tilde{\nu}_1^I$  denotes the lightest CP-odd sneutrino particle.

CP-even heavy $h_s$ scenario: $\ln Z = -31.8 \pm 0.02$			
Annihilation characteristics		Percent	
Coannihilation	$\tilde{\nu}_1 \chi_2 \rightarrow XY$	85%	86.1%
	$\tilde{\nu}_1 \tilde{\nu}_1^I \rightarrow Y_1 Y_2$	1.0%	
	$\tilde{\nu}_1^I \tilde{\nu}_1^I \rightarrow Y_3 Y_4$	0.1%	
Secluded DM sector	$\tilde{\nu}_1 \tilde{\nu}_1 \rightarrow A_s A_s$	2.9%	7.1%
	$\tilde{\nu}_1 \tilde{\nu}_1 \rightarrow h_s h_s$	0.2%	
	$\tilde{\nu}_1 \tilde{\nu}_1 \rightarrow \nu_4 \nu_4$	4.0%	
Higgs portal	$\tilde{\nu}_1 \tilde{\nu}_1 \rightarrow h h_s$	0.1%	6.8%
	$\tilde{\nu}_1 \tilde{\nu}_1 \rightarrow h h$	0.2%	
	$\tilde{\nu}_1 \tilde{\nu}_1 \rightarrow t \bar{t}$	0.1%	
	$\tilde{\nu}_1 \tilde{\nu}_1 \rightarrow b \bar{b}$	1.1%	
	$\tilde{\nu}_1 \tilde{\nu}_1 \rightarrow W^+ W^-$	5.4%	

**Table 4:** The annihilation mechanisms and channels in CP-even heavy  $h_s$  scenario, where  $\chi_2 \equiv \{\tilde{\chi}_1^0, \tilde{\chi}_1^\pm\}$ ,  $Y_1 Y_2 \equiv \{h A_s, \nu_4 \nu_4, \nu_5 \nu_5\}$ , and  $Y_3 Y_4 \equiv \{A_s A_s, W^+ W^-, \nu_4 \nu_4\}$ .

- ① Singlet-dominated particles,  $\tilde{\nu}_1^0$ ,  $h_s$ ,  $A_s$ , and  $\nu_h$ , may form a secluded DM sector.  $\lambda_\nu$ ,  $\kappa$  and  $v_s$  play an crucial role in determining the abundance.
- ② Due to limited theoretical framework, DM prefers to co-annihilate with Higgsino-dominated  $\tilde{\chi}_1^0$  to obtain the measured abundance.
- ③ Since constraints from DM experiments on electroweakinos and sleptons are weak, the theory can readily explain the muon g-2. For most cases, the Higgsino-dominated  $\tilde{\chi}_1^0$  appears as missing track at the LHC, and can be treated as an effective DM candidate. In this case, constraints from the LHC search for SUSY is weak.

## The status of Type-I NMSSM:

Model	MSSM	Z <sub>3</sub> -NMSSM		GNMSSM	Type-I NMSSM	B-L NMSSM
DM candidate	Bino	Bino	Singlino	Singlino	Sneutrino	Bino, Singlino, BLino, Bileptino, and sneutrino
DM physics	×	×	×	✓	✓	✓
LHC and $\Delta a_\mu$	×	×	×	✓	✓	✓
EWSB	×	×	✓	✓	✓	✓
Neutrino	×	×	×	×	✓	✓
Higgs mass	×	×	×	×	✓	✓

## Section Seven

### Example V: B-L NMSSM



## • Vector Superfields

SF	Spin $\frac{1}{2}$	Spin 1	$SU(N)$	Coupling	Name
$\hat{B}$	$\lambda_{\hat{B}}$	$B$	$U(1)$	$g_1$	hypercharge
$\hat{W}$	$\lambda_{\hat{W}}$	$W$	$SU(2)$	$g_2$	left
$\hat{g}$	$\lambda_{\hat{g}}$	$g$	$SU(3)$	$g_3$	color
$\hat{B}'$	$\lambda_{\hat{B}'}$	$B'$	$U(1)$	$g_B$	$B - L$

## • Chiral Superfields

SF	Spin 0	Spin $\frac{1}{2}$	Generations	$(U(1) \otimes SU(2) \otimes SU(3) \otimes U(1))$
$\hat{q}$	$\tilde{q}$	$q$	3	$(\frac{1}{6}, \mathbf{2}, \mathbf{3}, \frac{1}{6})$
$\hat{l}$	$\tilde{l}$	$l$	3	$(-\frac{1}{2}, \mathbf{2}, \mathbf{1}, -\frac{1}{2})$
$\hat{H}_d$	$H_d$	$\tilde{H}_d$	1	$(-\frac{1}{2}, \mathbf{2}, \mathbf{1}, \mathbf{0})$
$\hat{H}_u$	$H_u$	$\tilde{H}_u$	1	$(\frac{1}{2}, \mathbf{2}, \mathbf{1}, \mathbf{0})$
$\hat{d}$	$\tilde{d}_R^*$	$d_R^*$	3	$(\frac{1}{3}, \mathbf{1}, \mathbf{\bar{3}}, -\frac{1}{6})$
$\hat{u}$	$\tilde{u}_R^*$	$u_R^*$	3	$(-\frac{2}{3}, \mathbf{1}, \mathbf{\bar{3}}, -\frac{1}{6})$
$\hat{e}$	$\tilde{e}_R^*$	$e_R^*$	3	$(1, \mathbf{1}, \mathbf{1}, \frac{1}{2})$
$\hat{S}$	$S$	$\tilde{S}$	1	$(0, \mathbf{1}, \mathbf{1}, \mathbf{0})$
$\hat{\nu}$	$\tilde{\nu}_R^*$	$\nu_R^*$	3	$(0, \mathbf{1}, \mathbf{1}, \frac{1}{2})$
$\hat{\eta}_1$	$\eta_1$	$\tilde{\eta}_1$	1	$(0, \mathbf{1}, \mathbf{1}, -1)$
$\hat{\eta}_2$	$\eta_2$	$\tilde{\eta}_2$	1	$(0, \mathbf{1}, \mathbf{1}, 1)$

- Superpotential

$$W_{\text{B-L}} = W_{\text{GNMSSM}} + Y_\nu \hat{\nu} \hat{l} \hat{H}_u + Y_x \hat{\nu} \hat{\eta}_1 \hat{\nu} - \lambda_\eta \hat{s} \hat{\eta}_1 \hat{\eta}_2 + \mu_\eta \hat{\eta}_1 \hat{\eta}_2$$

- Naturally provide a seesaw mechanism for neutrino mass and mixing and the Bileptino mass.
- R-parity is related to gauge symmetry.

- Mass matrix for Neutralinos in the basis  $(\lambda_{\tilde{B}}, \tilde{W}^0, \tilde{H}_d^0, \tilde{H}_u^0, \lambda_{\tilde{B}'}, \tilde{\eta}_1, \tilde{\eta}_2, \tilde{S})$

$$\begin{pmatrix} M_1 & 0 & -\frac{1}{2}g_1 v_d & \frac{1}{2}g_1 v_u & M_{BB'} & -g_{BY} v_\eta & g_{BY} v_{\tilde{\eta}} & 0 \\ 0 & M_2 & \frac{1}{2}g_2 v_d & -\frac{1}{2}g_2 v_u & 0 & 0 & 0 & 0 \\ -\frac{1}{2}g_1 v_d & \frac{1}{2}g_2 v_d & 0 & m_{\tilde{H}_d^0 \tilde{H}_u^0} & -\frac{1}{2}g_{YB} v_d & 0 & 0 & -\frac{1}{\sqrt{2}}\lambda v_u \\ \frac{1}{2}g_1 v_u & -\frac{1}{2}g_2 v_u & m_{\tilde{H}_d^0 \tilde{H}_u^0} & 0 & \frac{1}{2}g_{YB} v_u & 0 & 0 & -\frac{1}{\sqrt{2}}\lambda v_d \\ M_{BB'} & 0 & -\frac{1}{2}g_{YB} v_d & \frac{1}{2}g_{YB} v_u & M_{BL} & -g_B v_\eta & g_B v_{\tilde{\eta}} & 0 \\ -g_{BY} v_\eta & 0 & 0 & 0 & -g_B v_\eta & 0 & m_{\tilde{\eta}_2 \tilde{\eta}_1} & m_{\tilde{S} \tilde{\eta}_1} \\ g_{BY} v_{\tilde{\eta}} & 0 & 0 & 0 & g_B v_{\tilde{\eta}} & m_{\tilde{\eta}_1 \tilde{\eta}_2} & 0 & m_{\tilde{S} \tilde{\eta}_2} \\ 0 & 0 & -\frac{1}{\sqrt{2}}\lambda v_u & -\frac{1}{\sqrt{2}}\lambda v_d & 0 & m_{\tilde{\eta}_1 \tilde{S}} & m_{\tilde{\eta}_2 \tilde{S}} & m_{\tilde{S} \tilde{S}} \end{pmatrix}$$

where

$$\begin{aligned} m_{\tilde{H}_u^0 \tilde{H}_d^0} &= -\frac{1}{\sqrt{2}}\lambda v_s - \mu, & m_{\tilde{\eta}_1 \tilde{\eta}_2} &= -\frac{1}{\sqrt{2}}\lambda_\eta v_s + \mu_\eta, \\ m_{\tilde{\eta}_1 \tilde{S}} &= -\frac{1}{\sqrt{2}}\lambda_\eta v_{\tilde{\eta}}, & m_{\tilde{\eta}_2 \tilde{S}} &= -\frac{1}{\sqrt{2}}\lambda_\eta v_\eta, & m_{\tilde{S} \tilde{S}} &= \sqrt{2}\kappa v_s + M_S. \end{aligned}$$

## Notes

Possible DM candidates:

Bino-, Singlino-, Blino-, Bilepton-dominated neutralino, and sneutrino.

Possible light particles: singlino-dominated Higgs, Bileptonic CP-even Higgs.

Singlet-dominated particles can naturally form a secluded DM sector.

## The status of B-L NMSSM:

Model	MSSM	Z <sub>3</sub> -NMSSM		GNMSSM	Type-I NMSSM	B-L NMSSM
DM candidate	Bino	Bino	Singlino	Singlino	Sneutrino	Bino, Singlino, BLino, Bileptino, and sneutrino
DM physics	×	×	×	✓	✓	✓
LHC and $\Delta a_\mu$	×	×	×	✓	✓	✓
EWSB	×	×	✓	✓	✓	✓
Neutrino	×	×	×	×	✓	✓
Higgs mass	×	×	×	×	✓	✓

# Part Eight

## Conclusions

# Conclusion about global fit of supersymmetric theories

- ① Experimental data provides many hints to fundamental physics.
- ② Global fit deepens greatly our understanding of new physics.
- ③ Economic supersymmetric theories are facing increasingly strong experimental restrictions, and more complex theory becomes favored to alleviate the constraints.
- ④ Some seeming independent problems may have a common physical origin. Well motivated theories should be explored in a more sophisticated way.

thanks!