

The Quantum Magic of Standard Model Parameters

Based on 2509.18251 and 2610.11121 with Qiaofeng Liu and Ian Low

Zhewei Yin (殷喆伟)



中国科学院高能物理研究所
Institute of High Energy Physics, Chinese Academy of Sciences

Oct 24, 2026

@ The International Conference on Fundamental Physics
from Quanta to Cosmos, Beijing

- Our framework for microscopic particle scattering: Quantum Field Theory (QFT)
- The final state of a scattering process naturally carries a lot of “quantumness”

For example: **entanglement** $\propto$ **cross section**

Low, **ZY**, 2405.08056, 2410.22414

See also Aoude, Elor, Remmen, Sumensari, 2402.16956; Zheng, Gu, 2610.10704

QI features: Property or principle?



In recent years, maximal or minimal entanglement has been found to single out unique theories

- Maximal entanglement $\rightarrow$ QED

Cervera-Lierta, Latorre, Rojo, Rottoli; Fedida, Serafini; ...

- Minimal entanglement $\rightarrow$ classical BH scattering, strong interactions, flavor structures in the SM, BSM theories, ...


Beane, Kaplan, Klco, Savage; Aoude, Chung, Huang, Machado, Tam;
Low, Mehen; Liu, Low, Mehen; Carena, Low, Wagner, Xiao;
Thaler, Trifinopoulos; Núñez, Cervera-Lierta, Latorre; ...



Quantum information properties of
particle scattering

? $\updownarrow$

Underlying principles of QFT,
e.g. symmetry



Quantum advantage

Entanglement alone is **not** enough to guarantee a quantum computational advantage

- Gottesman–Knill theorem: classical algorithms can efficiently simulate **stabilizer states** evolving through quantum circuits made of **Clifford gates**
- Clifford gates: the Hadamard gate H , the phase gate S , and the CNOT gate
 - Universal quantum computing: Clifford gates + the T gate

For a single qubit:

$$H = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}, \quad S = \begin{bmatrix} 1 & 0 \\ 0 & i \end{bmatrix},$$
$$T = \begin{bmatrix} 1 & 0 \\ 0 & e^{i\pi/4} \end{bmatrix}$$

For 2 qubits:

$$\text{CNOT}_{12} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

Magic, or non-stabilizerness: quantifies the quantum advantage over classical algorithms

- A computable measure: the stabilizer Rényi entropy (SRE)

Leone, Oliviero, Hamma, 2106.12587


Explored in the context of particle scattering:
nuclear physics, collider physics, neutrino physics, ...

Robin, Savage, 2024; White, White, 2024; Chernyshev, Robin, Savage, 2024; ...

Stabilizer states and magic

Stabilizer states: zero magic

- For a single qubit:
the 6 Pauli states



Credit: García-Álvarez, Ferraro, Ferrini, 1911.12346

- For 2 qubits: 60 states, with
zero or maximal entanglement

The α -SRE:

$$M_\alpha(|\psi\rangle) = \frac{1}{1-\alpha} \ln \sum_{\mathcal{O} \in \mathcal{W}(d)} \frac{1}{d} |\langle \psi | \mathcal{O} | \psi \rangle|^{2\alpha},$$

where $\mathcal{W}(d)$ is the Weyl–Heisenberg (WH) group

- For a single qubit: $\mathcal{W}(2) = \{\mathbf{1}, \sigma_x, \sigma_y, \sigma_z\}$
- For n -qubit systems: $\mathcal{W}(2)^{\otimes n}$

No optimization over states needed!

M_α quantifies, in a specific sense, the “distance”
from a state to the set of stabilizer states

- $M_2(|T\rangle) = \ln(3/2)$

Maximal magic for 2 qubits

What is the maximal SRE for a 2-qubit system?

- $M_2 \leq \ln(16/7) \approx 0.827$

Liu, Low, **ZY**, 2502.17550

- States saturating this bound: fiducial states for Weyl–Heisenberg mutually unbiased bases

Feng, Luo, *Commun.Theor.Phys.* 77 (2025) 1, 015102

- They are partially entangled: concurrence $1/2$ or $1/\sqrt{2}$

Quantum Sci. Technol. **11** (2026) 015035

Quantum Science and Technology

PAPER

Maximal magic for two-qubit states

Qiaofeng Liu¹, Ian Low^{1,2} and Zhewei Yin^{1,2,*} 


Magic in electroweak processes

Consider $e^-e^- \rightarrow e^-e^-$ and treat the spins of the two fermions as two qubits

- Treat θ_W as a free parameter


Average over the 60 stabilizer initial states...

$$\mathcal{M}_2(\theta) \equiv \frac{1}{60} \sum_{i=1}^{60} M_2(|\psi_s\rangle_i, \theta)$$



...as well as over the scattering angle

$$\langle \mathcal{M}_2 \rangle \equiv \int \frac{d\Omega}{4\pi} \mathcal{M}_2(\theta)$$



Liu, Low, ZY, 2509.18251


Magic for the weak mixing angle

The Standard Model value of s_W^2 tends to **minimize** the average magic

Liu, Low, **ZY**, 2509.18251

For charged leptons,
 $s_W^2 = 1/4 \Rightarrow$ the Z coupling is purely axial


- $e^-e^- \rightarrow e^-e^-$: t and u channels
- $e^-\mu^- \rightarrow e^-\mu^-$: t channel only



Magic in QCD processes

Consider color-singlet $q\bar{q} \rightarrow q\bar{q}$, treating the two fermion spins as two qubits

- Treat α_s/α as a free parameter
- Light quarks: $\alpha_s > \alpha$ is preferred





Liu, Low, **ZY**, 2610.11121

Magic for the strong coupling constant

$t\bar{t}$: the Standard Model value of α_s tends to **minimize** the average magic


Liu, Low, **ZY**, 2610.11121



Magic for the Higgs sector?

- Higgs exchange is essential for the SM magic to lie close to the minimum
- Removing the Higgs exchange channel, i.e. taking $m_h \rightarrow \infty$, significantly worsens the agreement

Liu, Low, **ZY**, 2610.11121



Implications: minimizing the resource for quantum computation as an underlying principle of fundamental physics?

- Why the SRE? Other measures of magic?

Shen, **ZY**, Zhang, *upcoming*

- The prescription for the quantum simulation, e.g. basis dependence?

Li, Low, Wang, **ZY**, 2608.19095

- Which processes determine which SM parameters? Quantitative predictions?
- The mechanism behind magic minimization?

Thank you!